

Influence of Glutamine and Vitamin E on the Performance and the Immune Responses of Broiler Chickens¹

■ Author(s)

Sakamoto MI²
Murakami AE³
Silveira TGV⁴
Fernandes JIM²
Oliveira CAL de²

- ¹ Part of the dissertation thesis of the first author presented in the Post-Graduation Program in Animal Science of Universidade Estadual de Maringá/UEM-PR, financed by pelo Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq.
- ² Post-Graduation Program in Animal Science of Universidade Estadual de Maringá/UEM-PR.
- ³ Department of Animal Science – Post-Graduation Program in Animal Science of Universidade Estadual de Maringá/UEM-PR.
- ⁴ Department of Clinical Analyses of Universidade Estadual de Maringá/UEM-PR.

■ Mail Address

Márcia Izumi Sakamoto
Rua Duílio Vasecchi, 204. Apto. 03
Jardim Petrópolis
13.634-304. Pirassununga, SP, Brazil

Email: mizumiss@yahoo.com.br
mizumiss@hotmail.com

■ Keywords

Antibodies, sheep red blood cells, cutaneous basophilic hypersensitivity.

ABSTRACT

This study aimed at evaluating the influence of Glutamine (Gln) and Vitamin E (VE) supplementation on the performance and immune response of broilers. A completely randomized experimental design with a 2 x 3 (VE x Gln) factorial arrangement was used. VE was supplemented at 10 and 500 mg/kg feed, with or without Gln (1%) addition, and two periods of supplementation in the starter diets (1-7 and 1-14 days of age), with five replicates of 50 birds each. The analyzed parameters were: live performance (weight gain, feed intake, and feed conversion ratio); relative weights of the spleen, bursa, and thymus; antibody titers (with sheep red blood cells suspension – SRBC) and cutaneous basophilic hypersensitivity (CBH). Data were submitted to the analysis of variance, and means were compared using the test of Tukey. Treatments did not influence ($P>0.05$) live performance parameters or antibody titers. VE reduced ($P=0.01$) CBH, with the level of 10 mg VE/kg allowing higher cell proliferation as compared to 500 mg VE/kg. As to lymphoid organs, only the spleen was affected ($P=0.035$) by Gln, which resulted in higher spleen relative weight when fed during the first week of age. Results showed that 10 mg VE/kg with Gln (1-7 days) promoted better immune responses.

INTRODUCTION

Action of vitamin E on the immune response

Vitamin E (VE) is the most widely known natural anti-oxidant, and alpha-tocopherol is its most active biological form for physiological functions, despite its limited efficiency as anti-oxidant. The delta and gamma forms are better anti-oxidants, but are less efficient to support animal growth and performance (Rutz & Lima, 1994). According to McDowell (1989), DL-alpha-tocopherol power is 1.1 IU/mg of Vitamin E.

VE donates electrons to free radicals, thereby making them stable. This prevents free radicals from binding to fatty acids, inhibiting oxidation reactions, and therefore maintaining cell membrane integrity (Rutz & Lima, 1994).

According to Klasing (1998), in addition of being the first line of body defenses against the action of free radicals, protecting the host cells, VE modulates the immune response. It also decreases the synthesis of prostaglandins, leukotrienes, and cytokines, which regulate the inflammatory response, thereby reducing damage caused to the tissues by the inflammatory process.

Chung & Boren (1999) evaluated the benefits of VE supplementation in broiler diets, and observed that birds fed 240 mg VE/kg improved feed conversion ratio in 2.3% as compared to the control group (33 mg de VE/kg). Broilers that received the higher VE level presented reductions of 34, 25, and 61% in carcass downgrading due to diseases, septicemia/



toxemia, and inflammatory processes, respectively, relative to the control group.

On the other hand, Erf *et al.* (1998) supplemented broiler diets with increasing VE levels (0, 17, 46 e 87 mg de dl- α -tocopherol/kg), did not find effect of VE supplementation on thymus and spleen B lymphocytes and macrophages percentage. However, independent of supplementation level, VE increased T-cell (CD4⁺ and CD8⁺) production as compared to birds that did not receive VE supplementation.

Boa-Amponsem *et al.* (2000) fed broiler diets containing 10 and 300 mg VE/kg, and challenged the birds with an intravenous solution of sheep red blood cells. There was no effect of VE supplementation on anti-SRBC antibody titers, but the heterophil/lymphocyte ratio increased, indicating that VE improved the phagocytic capacity of the immune system, protecting the birds against the invasion of pathogenic microorganisms.

Evaluating VE supplementation on (0, 10, 25, 50, 100, and 200 IU DI- α -tocopherol/kg) on the immune response of broilers, Leshchinsky & Klasing (2001) observed that VE levels between 25 and 50 IU/kg (22.73 and 45.45 mg/kg) promoted higher antibody titers 7 days after the inoculation of SRBC as compared to higher levels of VE dietary addition.

Glutamine action on the immune response

Glutamine (Gln) is a free, neutral, non-essential amino acid, which found in higher levels in muscles and plasma, in concentrations representing approximately 50 to 80% of the total free amino acids content in the body. As it contains in its structure two nitrogen groups that can be mobilized, Gln may be a vehicle for tissue nitrogen exchange, and may play an essential role in several important metabolic pathways (Marliss *et al.*, 1971; Smith, 1990).

Gln is recognized as a crucial energy substrate for rapidly dividing cells, and may act on the humoral immune response, that is, in certain sites of mucosa membranes, such as the respiratory and gastrointestinal tracts, with increase in the number of lymph nodes in mammals (Newsholme, 2001).

Yi *et al.* (2005) observed the importance of Gln supplementation (1%) in the diet of broilers up to 28 days of age, which presented better performance (weight gain, feed efficiency, and viability) than the non-supplemented birds. However, these results are not in agreement with the study of Maiorka *et al.* (2000), who did not find any effect of Gln on the performance of broilers supplemented at the same levels and during the same period.

Although the immunomodulation mechanism of several dietary ingredients is still being researched, the outstanding importance of dietary influence on the immune response should not be underestimated in terms of the general resistance of poultry against infections.

The immune system has one of the most complex molecular and cell interactions in biology, encompassing the cell system (T cells) and the humoral system (B cells), and the lymphocyte is the primary cell in both systems. Antibodies are produced in the secondary lymphoid tissues, which include, in addition to the spleen and the lymph nodes, the bone marrow, the cecal tonsils, and the lymphoid tissue distributed throughout the body, particularly in the respiratory, digestive, and urogenital tracts (Tizard, 1985).

At hatching, the immune system of birds is already partially developed, and the primary organs – thymus and bursa – are present and populated with lymphoid cells. However, the secondary organs, such as spleen, cecal tonsils, Meckel's diverticulum, and lymphoid tissues scattered in the digestive and respiratory tract are still incomplete (Dibner & Richards, 2004).

Intensive genetic selection for weight gain and feed efficiency caused poultry to be less competent in some aspects of the immune response. When commercial genetic lines of broilers were compared, it was observed that 1991 strains were less efficient in antibody production as compared to 1957 strains. Interestingly, genetic manipulation did not change natural immune response (Qureshi & Havestein, 1994).

Intensive production systems increase the risk of dissemination of infectious diseases, demanding the adoption of measures to preserve bird health. Production practices, in addition of increasing pathogen load, also caused changes in the environment, both in terms of climate and behavior, to which broilers were not adapted (Siegel, 1985).

The ban of some feed additives used to improve broiler performances and profitability drove the study of other ingredients that could allow optimal nutrient utilization, favoring the full expression of the genetic potential (Perry, 1995; Rosen, 1996).

Aiming at contributing to broiler nutrition research, and considering the lack of information in literature as to these two nutrients – glutamine and vitamin E – this experiment evaluated the influence of Gln and VE supplementation on the performance and on the intensity of the immune response of broilers.



MATERIAL AND METHODS

The experiment was carried out at the Poultry Production Sector of Iguatemi Experimental Farm of the State University of Maringá, and at the Animal Nutrition (LANA/DZO) and Immunogenetics (DAC) Laboratories of that university. The experimental procedures were approved by the Committee on Animal Experimentation Ethics – CEEA/UEM.

A total number of 1,500 day-old male Cobb-Vantress® broilers was used. A completely randomized experimental design with a 2 x 3 (VE vs. Gln) factorial arrangement was applied. VE levels of 10 and 500 mg (11 and 550 IU/kg, respectively) of DL- α -tocopherol/kg were added to the feed, with or without Gln (1%) added during two starter periods (from 1 to 7 days and 1 to 14 days of age), with a total number of six treatments, with five replicates of 50 birds per experimental unit. During the grower state (from 22 to 41 days of age), treatments consisted only of the two different VE levels. The vitamin and mineral premix used in the feeds did not contain VE. Diets contained equal nutrient levels, and were based on corn and soybean meal, being formulated to supply nutritional requirements according to Rostagno *et al.* (2000). The experimental diets are shown in Tables 1 and 2.

Table 1 – Percentage and calculated¹ composition of starter broiler feeds (1 to 21 days of age).

Ingredients	Vitamin E level (mg/kg feed)			
	10		500	
	no Gln	with Gln (1%)	no Gln	with Gln (1%)
Corn grain	55.86	55.86	55.77	55.77
Soybean meal 45%	36.34	36.34	36.35	36.35
Dicalcium phosphate	1.82	1.82	1.82	1.82
Limestone	1.08	1.08	1.08	1.08
Soybean oil	2.85	2.85	2.88	2.88
Salt	0.40	0.40	0.40	0.40
DL-methionine (99%)	0.23	0.23	0.23	0.23
Mineral-vitamin premix ²	0.40	0.40	0.40	0.40
Vitamin E ³	0.001	0.001	0.05	0.05
L-Glutamine ⁴	-	1.00	-	1.00
Inert material ⁵	1.00	-	1.00	-
BHT ⁶	0.01	0.01	0.01	0.01
Total	100.00	100.00	100.00	100.00

1 – Calculated requirements: 3,000 kcal ME/kg; 21.50 % CP; 1.00% Ca; 0.45% Avail. P; 0.90% met+cys; 1.15% Lys; 0.20% Na; 0.27% Cl; 0.84% K. 2 – Nutritional levels per kg product: vitamin A, 12,500 IU; vitamin D3, 2,125 IU; vitamin K3, 650 mg; thiamin, 2,250 mg; riboflavin, 5,100 mg; pyridoxine, 2,475 mg; cyanocobalamine, 14,375 mcg; calcium pantothenate, 14,500 mg; niacin, 27,425 mg; folic acid, 0,800 mg; choline, 187,500 mg; zinc, 90,740 mg; iron, 82,250 mg; copper, 18,120 mg; manganese, 79,137 mg; iodine, 1,239 mg; selenium, 0,363 mg; cobalt, 2,498 mg; gentian violet, 3,125 mg; BHT, 1,250 mg. 3 – Vitamin E (DL- α -tocopherol): 60 %. 4 – Ajinomoto Interamericana: Analytical results - Transmittance 98%; Chloride (Cl) 0.020%; Ammonium (NH₄) 0.10%; Sulfate (SO₄) 0.020%; Iron (Fe) 10ppm; Heavy metals (Pb) 10 ppm; Arsenic (As₂O₃) 1 ppm; Loss on drying 0.01%; Assay 100.3%; pH 5.6. 5- Washed sand, replacing used glutamine. 6- Butyl Hydroxi Toluene (Antioxidant).

Table 2 - Percentage and calculated¹ composition of grower broiler feeds (21 to 41 days of age).

Ingredients	Vitamin E level (mg/kg feed)	
	10	500
Corn grain	64.89	64.78
Soybean meal 45%	29.19	29.22
Dicalcium phosphate	1.58	1.58
Limestone	1.02	1.02
Soybean oil	2.34	2.38
Salt	0.32	0.32
DL-methionine (99%)	0.19	0.19
L-Lysine HCl (78 %)	0.03	0.03
Mineral-vitamin premix ²	0.40	0.40
Vitamin E ³	0.001	0.05
BHT ⁴	0.01	0.01
Total	100.00	100.00

1 – Calculated requirements: 3,100 kcal ME/kg; 19 % CP; 0.90% Ca; 0.40% Avail. P; 0.80% met+cys; 1.00% Lys; 0.17% Na; 0.23% Cl; 0.73% K. 2 – Nutritional levels per kg product: vitamin A, 12,500 IU; vitamin D3, 2,125 IU; vitamin K3, 650 mg; thiamin, 2,250 mg; riboflavin, 5,100 mg; pyridoxine, 2,475 mg; cyanocobalamine, 14,375 mcg; calcium pantothenate, 14,500 mg; niacin, 27,425 mg; folic acid, 0,800 mg; choline, 187,500 mg; zinc, 90,740 mg; iron, 82,250 mg; copper, 18,120 mg; manganese, 79,137 mg; iodine, 1,239 mg; selenium, 0,363 mg; cobalt, 2,498 mg; gentian violet, 3,125 mg; BHT, 1,250 mg. 3 – Vitamin E (DL- α -tocopherol): 60 %. 4 – Butyl Hydroxi Toluene (Antioxidant).

Feeds were offered according to the established stages. Birds were weighed in the beginning and at the end of each period in order to determine performance (average feed intake, average weight gain, and feed conversion ratio). Mortality, as well as feed residues, was duly recorded to determine true feed intake.

Litter consisted of wood shavings, and was used for 5th time in order to increase challenge during grow-out. House temperature was daily recorded using a manual thermometer, and the maximum and minimum average temperatures were 30° C and 22° C, respectively. Average relative humidity was 30%. Mortality recorded during the entire experimental period was 1.50%.

Antibody titers (Humoral response)

On days 14 and 24, two birds were randomly removed from each replicate, and were duly identified with a wing band. These birds were inoculated with 0.5 ml of a suspension of sheep red blood cells (SRBC) diluted in PBS (phosphate buffered saline solution) at 0.5%. Inoculation was intramuscular in five equally distant points in the bird's breast, injecting 0.1 ml per point. On days 14 and 35 (days 0 and 21 after the first inoculation), blood was collected from the brachial vein, 3ml/bird. Blood was centrifuged at 3,000 rpm for 10 min, and the obtained serum was stored in duly identified 1-ml Eppendorf tubes in a freezer at -20°C until analyses. Before analyses, the serum were



inactivated in water batch at 56° C for 30 min, using as diluent buffer PBS with 1% bovine serum albumin (BSA/Fort Dodge®-22%). Humoral specific immune response was assessed by determination of anti-SRBC antibody titer, using the technique of simple hemagglutination, as described by Wegmann & Smithies (1965).

Two other birds per replicate were submitted to the same procedure, but these birds were injected with saline solutions to be used as negative controls. Titers were expressed as log₂ of the highest dilution observed in agglutination.

Cutaneous Basophilic Hypersensitivity – CBH (Cell Response)

At 36 days of age, two birds per replicate were used to assess late unspecific immune response. Each bird was identified with a band in the left wing, and submitted to intradermal inoculation in the inter-digital space between the 3rd and the 4th finger of the right foot with 100 µg of phytohemagglutinin-P (PHA-P – Gibco®), diluted in sterile saline solution, at a dose of 0.10 ml. An equivalent volume of sterile saline solution was injected in the same inter-digital space on the left foot for negative control.

Skin thickening of the inter-digital space was measured in millimeters in both feet using a manual pachymeter immediately before injection (time zero), 6, 12, and 24 hours after PHA-P injection. The response to cutaneous hypersensitivity test was calculated as follows (Corrier & DeLoach, 1990):

CBH= PHA-P response (right foot) – control response (left foot)

Where:

PHA-P response = post-injection thickening (right foot) – thickening at time zero (right foot)

Control response = post-injection thickening (left foot) – thickening at time zero (left foot)

Relative weight of lymphoid organs

At 41 days of age, two birds per replicate were sacrificed by electric stunning, followed by bleeding, in order to collect the spleen, the thymus, and the bursa. Relative weight of each organ was calculated as follows:

$$\text{Relative weight} = (\text{organ weight/live body weight}) \times 100$$

Statistical analysis

Data obtained for each parameter were submitted to analysis of variance, and means were tested ($P \leq 0.05$), according to their distributions, using the GLM procedure of SAS software (2000). Polynomial regression was applied to CBH (cell immunity) data as a function of measured hours.

The following statistical model was applied:

$$Y_{ijk} = m + V_i + G_j + VG_{ij} + e_{ijk}$$

Where:

Y_{ijk} = observation of the i^{th} replicate submitted to treatment of level V_i , and G_j ;

m = general mean of all observations;

V_i = effect of the i level of VE supplementation, $i = 1, 2$ (1= 10; 2= 500 mg VE/kg);

G_j = effect of j period with Gln (1%), $j = 1, 2, 3$ (1= exempt; 2= from 1 to 7 days; 3= from 1 to 14 days of age);

VG_{ij} = effect of VE vs. Gln interaction;

e_{ijk} = experimental error.

RESULTS AND DISCUSSION

Average results of performance characteristics are presented in Table 3. There was no effect ($P > 0.05$) of Gln and VE association on broiler performance during the studied rearing periods; however, low mortality percentage was observed during the entire experimental period. Reused wood shavings litter did not pose higher challenge. Our results are consistent with those of Maiorka *et al.* (2000), who did not observe any effect of Gln supplementation on broiler performance, and with those of Konjufca *et al.* (2004), who evaluated different VE levels, but no effects on broiler body weight were found. On the other hand, Yi *et al.* (2005) reported better feed efficiency, weight gain, and viability of broilers fed 1% Gln.

In terms of lymphoid organ relative weight, only the spleen was affected ($P = 0.035$) by Gln feeding during the starter period. Birds fed Gln during the first week of age presented higher relative spleen weight, as shown in Table 4. The rate of Gln utilization is high in isolated cells of the immune system, such as lymphocytes, macrophages, and neutrophils. In addition, Gln is important for lymphocyte proliferations, cytokine production, as well as for the activities of phagocytosis and secretion by the macrophages (Newsholme, 2001). These reasons could therefore



Table 3 – Mean values of performance traits (final weight, weight gain, individual feed intake, feed conversion ratio) and mortality percentage of 1 to 41-day-old broilers.

Treatment	Vitamin E (mg/kg)		Glutamine			MEAN	CV (%) ¹
	10	500	0	1 - 7 days	1 - 14 days		
Period (days)			Final Weight (g)				
1 to 7	168.267	165.574	164.680	168.382	167.700	166.921	3.25
1 to 14	442.994	437.757	437.241	441.469	442.416	440.376	3.13
1 to 21	875.564	879.388	875.863	887.565	869.000	877.476	3.51
1 to 41	2612.385	2619.838	2610.463	2613.215	2624.657	2616.112	2.49
			Weight Gain (g)				
1 to 7	123.530	120.798	120.084	123.654	122.756	122.165	4.24
1 to 14	398.258	392.981	392.645	396.741	397.472	395.619	3.42
1 to 21	830.828	834.612	831.267	842.837	824.056	832.720	3.68
1 to 41	2567.649	2575.062	2565.867	2568.487	2579.713	2571.356	2.53
			Feed Intake (g/bird)				
1 to 7	139.520	136.840	136.520	139.660	138.360	138.180	3.52
1 to 14	507.133	500.951	500.960	505.077	506.089	504.042	2.55
1 to 21	1154.445	1147.962	1145.577	1155.748	1152.285	1151.203	2.39
1 to 41	4464.532	4484.482	4481.110	4463.683	4478.727	4474.506	2.20
			Feed Conversion Ratio (g/g)				
1 to 7	1.129	1.133	1.137	1.129	1.127	1.131	1.71
1 to 14	1.273	1.275	1.276	1.273	1.273	1.274	1.37
1 to 21	1.390	1.376	1.379	1.371	1.399	1.383	1.86
1 to 41	1.739	1.741	1.747	1.738	1.736	1.740	1.46
			Total Cumulative Mortality (%)				
1 to 7			0.13				
1 to 14			0.34				
1 to 21			0.49				
1 to 41			1.50				

(P>0.05). 1 - CV (%) of treatment mean.

Table 4 – Mean values of relative weights (% live body weight) of spleen, bursa, thymus, and bursa size¹ 41-day-old broilers.

	Spleen	Bursa	Thymus	Bursometer ¹
Vitamin E (mg/kg)				
10	0.126	0.063	0.475	4.233
500	0.129	0.060	0.479	4.133
P-value	0.418	0.515	0.868	0.383
Glutamine (1%)				
1- No Glutamine	0.131 ab	0.066	0.477	4.200
2- 1 to 7 days	0.134 a	0.061	0.460	4.150
3- 1 to 14 days	0.121 b	0.057	0.492	4.200
P-value	* 0.035	0.182	0.475	0.918
VE*Gln (P-value)	0.383	0.625	0.692	0.467
CV (%)	11.92	23.82	16.08	10.54

* Different letters in the same column are significantly different by the test of Tukey (P≤0.05). 1 - Mean bursa size values as measured by the ruler Solvay-Bursine[®]-2.

justify the increase in spleen relative weight observed in the present experiment.

Vitamin E did not influence (P>0.05) the relative weight of the studied organs, which is in agreement with Gore & Qureshi (1997), who did not observe any difference in the relative weight of the spleen and the bursa of 35-day-old broilers that had received 10 IU vitamin E as embryos on the 18th day of incubation. Konjufca *et al.* (2004), evaluating VE supplementation as DL- α -tocopherol, did not find differences in organ relative weight, except for the spleen, which presented an increase in the 7th week of age of broilers fed 110

and 220 mg VE/kg as compared to control birds fed 16 mg VE/kg (0.14, 0.18, and 0.10, respectively).

The studied VE levels presented similar behavior as to cell immune response (cutaneous basophilic hypersensitivity – CBH). CBH response was decreasingly detected as measured times decreased. At 12 hours after inoculation, the level of 10 mg VE/kg promoted higher cell proliferation as compared to the 500 mg VE/kg level. This effect was no longer observed 24 hours after inoculation, as seen in Table 5. VE supplementation higher than the NRC (1994) recommendations had an inhibitory effect on T-helper cells, as also observed by Boa-Amponsem *et al.* (2000), who found a higher CBH level in birds treated with 10 IU VE/kg ($x=0.40 \pm 0.03$ mm) as compared to 300 IU VE/kg ($x=0.30 \pm 0.03$ mm) 24 hours after inoculation.

Using a single PHA-P inoculation, CBH response provides pre-immunization information in a short period of time. This response to a cutaneous test allows the assessment of the immunocompetence of T-helper cells in newly-hatched birds. Corrier & DeLoach (1990) evaluated the effect of different PHA-P doses (100 and 200 μ g) on cutaneous hypersensitivity of broilers in the starter period (up to 14 days of age), and observed the presence of T-helper cells already in 3-day-old birds, which skin thickness was first observed 6 hours post-inoculation, reached its maximum level 12 hours, and



Table 5 – Mean CBH values¹ (mm) as a function of vitamin E levels of 36-day-old broilers².

Vitamin E	Measurement time (hours)			CV (%)
	6	12	24	
10 ³	0.577 ± 0.025	0.524 ± 0.021 a	0.299 ± 0.018	24.18
500 ⁴	0.552 ± 0.027	0.483 ± 0.023 b	0.250 ± 0.014	25.17

1- CBH= (thickening, PHA-P / right foot) – (thickening, control-saline solution / left foot). 2- Mean ± standard error. 3- CBH = 0,6900-0,0159524*X (R²= 0,53); - CBH = 0,6683- 0,0171429*X (R²= 0,59); * Different letters in the same column are significantly different by the test of Tukey (P≤0.05)

Table 6 – Mean values of antibody titers as measured in 35-day-old broilers fed diets supplemented with Glutamine and Vitamin E.

	Vitamin E (mg/kg)		0	Glutamine (1%)		CV (%)
	10	500		1 - 7 days	1 - 14 days	
Antibody titer ¹	2.733	2.400	2.500	2.650	2.550	25.05
Standard error	0.348	0.320	0.316	0.428	0.496	

1 - Values expressed as Log₂ of the highest dilution observed in Agglutination (P>0.05).

decreased 24 hours post-inoculation. Conversely, Leshchinsky & Klasing (2001) did not find any effect of VE supplementation on CBH of broilers.

In terms of humoral immune response, average antibody titers are presented in Table 6. There was no effect (P>0.05) of the treatments on antibody titers of birds at 35 days of age. However, birds treated with 10 mg VE/kg and Gln during the first week of age produced more antibodies.

Gore & Qureshi (1997) injected VE as DL- α -tocopherol in turkey embryos at 18 days of incubation, and observed that birds receiving 10 IU VE/kg presented higher antibody titers as compared with those that did not receive VE (4.2 *vs.* 3.0) at 14 days after SRBC inoculation. This suggests that VE may have an immunomodulator effect, increasing the resistance to diseases. A possible mechanism of this enhanced immune function may be a down regulation of the biosynthesis of prostaglandins, which inhibit several immunity parameters.

The results of the present experiment are similar to those observed by Boa-Amponsem *et al.* (2000), who did not find significant effect of vitamin E (levels of 10 and 300 IU/kg) on average antibody titers of broilers inoculated with SRBC, as measured 20 days after inoculation.

Yang *et al.* (2000) studied chickens with high and low antibody production and VE (10 and 300 IU/kg), and verified a decrease in antibody titers in chickens with low genetic potential fed the high VE level at 6 and 10 days after inoculation with SRBC (2.4 *vs.* 1.5 on day 6; and 1.9 *vs.* 1.2 on day 10, for 10 and 300 IU VE/kg, respectively).

On the other hand, Leshchinsky & Klasing (2001) found an increase in the antibody titers of broilers supplemented with 50 IU VE/kg (x= 3.35 at 7 days after inoculation), and concluded that moderate VE levels

(25 to 50 IU/kg) promoted better immunomodulation than high VE levels (100 to 200 IU/kg), which correspond to the VE levels needed for the inhibition of lipid peroxidation, and for the protection of liver mitochondria against oxidative stress.

CONCLUSIONS

The association of 10 mg vitamin E/kg (11 IU/kg) with Glutamine (during the first 7 days of age) in the feed of broilers promoted better immune response, but did not influence broiler performance.

REFERENCES

- Boa-Amponsem K, Price SEH, Picard M, Geraert PA, Siegel PB. Vitamin E and immune responses of broiler pureline chickens. *Poultry Science* 2000; 79:466-470.
- Chung TK, Boren B. Vitamin E use in commercial flocks examined. *Feedstuffs* 1999; 6:11-14.
- Corrier DE, DeLoach JR. Evaluation of cell-mediated cutaneous basophil hypersensitivity in young chickens by an interdigital skin test. *Poultry Science* 1990; 69:403-408.
- Dibner JJ, Richards JD. The digestive system: challenges and opportunities. *Journal Applied Poultry Research* 2004; 13:86-93.
- Erf GF, Bottje WG, Bersi TK, Headrick MD, Fritts CA. Effects of dietary vitamin E on the immune system in broilers: altered proportions of CD4 T cells in the thymus and spleen. *Poultry Science* 1998; 77:529-537.
- Gore AB, Qureshi MA. Enhancement of humoral and cellular immunity by vitamin E after embryonic exposure. *Poultry Science* 1997; 76:984-991.
- Klasing KC. Nutritional modulation of resistance to infectious diseases. *Poultry Science* 1998; 77:1119-1125.
- Konjufca VK, Bottje WG, Bersi TK, Erf GF. Influence of dietary vitamin



E on phagocytic functions of macrophages in broilers. *Poultry Science* 2004; 83:1530-1534.

Leshchinsky TV, Klasing KC. Relationship between the level of dietary vitamin E and the immune response of broiler chickens. *Poultry Science* 2001; 80:1590-1599.

Maiorka A, Silva AVF, Santin E, Borges SA, Boleli IC, Macari M. Influência da suplementação de glutamina sobre o desempenho e o desenvolvimento de vilos e criptas do intestino delgado de frangos. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia* 2000; 52(5):487-490.

Marliss EB, Aoki TT, Pozefsky T, Most AS, Cahill Jr GF. Muscle and splanchnic glutamine and glutamate metabolism in post absorptive and starved man. *Journal of Clinical Investigation* 1971; 50:814-817.

McDowell LR. Vitamins in animal nutrition: comparative aspects to human nutrition. London: Academic Press; 1989. p. 93-131.

National Research Council - NRC. Nutrient requirements of domestic animals. 9th ed. Washington, DC; 1994.

Newsholme P. Why is L-Glutamine metabolism important to cells of the immune system in health, postinjury, surgery or infection? *Journal of Nutrition* 2001; 131:2515S-2522S.

Perry FG. Biotechnology in animal feeds and animal feeding: an overview. In: Wallace RJ, Cheson A. *Biotechnology in animal feeds and animal feeding*. Local: Editora; 1995. p.1-15.

Qureshi MA, Havestein GB. A comparison of the immune performance of a 1991 commercial broiler with a 1957 randombred strain when fed "typical" 1957 and 1991 broiler diets. *Poultry Science* 1994; 73:1805-1812.

Rosen GD. Feed additive nomenclature. *World's Poultry Science Journal* 1996; 52:53-57.

Rostagno HS, Albino LFT, Donzele JL, Gomes PC, Ferreira AS, Oliveira RF, Lopes DC. Tabelas brasileiras para aves e suínos: composição de alimentos e exigências nutricionais. Viçosa: Editora Imprensa Universitária; 2000.

Rutz R, Lima GJMM. Uso de antioxidantes em rações e subprodutos. In: Conferência Apinco de Ciência e Tecnologia Avícolas; 1994; Santos, São Paulo. Brasil. Anais... Santos: Facta; 1994. p.73-84.

SAS INSTITUTE. SAS user's guide, statistics, versão 8.1. 4th ed. Cary, 2000.

Siegel HS. Immunological responses as indicators of stress. *World's Poultry Science Journal* 1985; 41:36-44.

Smith RJ. Glutamine metabolism and its physiologic importance. *Journal of Parenter Enteral Nutrition* 1990; 14:405-445.

Tizard I. Introdução à imunologia Veterinária. 2.ed. São Paulo: Editora Roca; 1985.

Wegmann TG, Smithies O. A simple hemagglutination system

requiring small amounts of red blood cells and antibodies. *Transfusion* 1965; 6:67-73.

Yang N, Larsen CT, Dunnington EA, Geraert PA, Picard M, Siegel PB. Immune competence of chicks from two lines divergently selected for antibody response to Sheep Red Blood Cells as affected by supplemental vitamin E. *Poultry Science* 2000; 79:799-803.

Yi GF, Allee GL, Knight CD, Dibner JJ. Impact of glutamine and oasis hatchling supplement on growth performance, small intestinal morphology, and immune response of broilers vaccinated and challenged with *Eimeria maxima*. *Poultry Science* 2005; 84:283-293.