



Analyzing Growth Curves of Turkeys Reared in Different Breeding Systems (Intensive and Free-Range) with some Nonlinear Models

■ Author(s)

Sogut B¹
Celik S¹
Ayasan T^{II}
Inci H¹

¹ Department of Animal Science, Faculty of Agriculture, Bingol University, Turkey

^{II} Eastern Mediterranean Agricultural Research Institute

■ Mail Address

Corresponding author e-mail address
Dr. Bunyamin Sogut
Department of Animal Science, Faculty of Agriculture, Bingol University Bingol/
TURKEY.
Email: bunyaminsogut@hotmail.com

■ Keywords

Breeding system, growth models, large white turkey, live weight.

ABSTRACT

The objective of this study was to identify the growth curve of live weight of turkeys reared in different systems (intensive and free-range) with the best mathematical model. To achieve this goal, the live weight of male and female turkeys (Large White) reared for 0-18 weeks were used. Logistic, Gompertz, Von Bertalanffy, and Gauss were used to determine the best model for the turkeys. In comparison of the models, values of Coefficient of Determination (R^2), Mean Squares of Error (MSE) and Model Efficiency (ME) were used. In Von Bertalanffy model, the coefficient of determinations for males and females were found as 0.9974 and 0.99915 in intensive system and 0.9794 and 0.9804 in Free-Range system, respectively.

As a result of this study, because the highest R^2 and the lowest MSE were observed in Von Bertalanffy model, it was the best among the models to identify growth curve of the turkeys.

INTRODUCTION

Growth is the change in weight and body sizes of the living organism, and it is an economically important feature in livestock breeding (Alpan and Arpacık, 1998; Akçapınar and Özbeyaz, 1999). Exposure of the economic importance of the features about live weight and growth rate has aroused of scientist's and grower's interest, and it has increased their interest in analyzing the weight-age relationship (Ngeno *et al.* 2010; Rizzi *et al.* 2013; Eleroglu *et al.* 2014).

The growth of the animals is showed up as a result of complex metabolic activities thus, researchers have tried to explain numerically the growth characteristics by improving various mathematical models to express this complex biologic situation better. Obtaining the expected performance of the animals at mature ages is related to these animal's growth and development. The changes related to the age in the growth of an animal is called the growth curve (Akbaş and colleagues, 1999; Malhado *et al.*, 2008; Agudelo-Gomez, 2009; Bayram and Akbulut, 2009; Prestes, 2012). Briefly, the changes shown in living growth and body sizes, based on the age, can be defined as growth curve (Goonewardene *et al.* 198; Kocabaş *et al.*, 1997). In the process of time that growth ends and the animal reaches the weight of mature living, biologic parameters which have an important place in explaining growth physically can be guessed (Behr *et al.*, 2001). Şengül and Kiraz (2005) have examined the growth curves of Large White turkeys, and have determined that Gompertz's, Logistic's, Richards's, and Morgan-Mercer-Flodin's models, gave nearly the same coefficient of determinations (99%).

In livestock-breeding, growth curves are used in subjects to identify the best slaughtering age, to get information about the general state



of health of the living, to identify the age that it can be used in breeding, to identify the age of sexual maturity (Doğan, 2003), to investigate the effects of selection on growth curve parameters, etc. (Bilgin *et al.* 2004; Tariq *et al.* 2011; Tariq *et al.* 2013).

The growth orbit measured as body mass and body weight is defined with proper mathematical methods relevant to growth curves, especially of poultry (Aggrey, 2002; Reddish and Lilburn, 2004; Norris *et al.*, 2007).

Growth is an important economic feature in broiler industry, and it can be defined as the increase of body size per unit, in time (Schulze *et al.*, 2001). The estimation of growth curve parameters can be of importance to the economy of production. Biological implications of the model parameters, and their relationship with other production features, provide a sound basis for developing a breeding strategy to modify the growth curve (Lambe *et al.* 2006; Abegaz *et al.* 2010; Ayied *et al.* 2011; Saghi *et al.* 2012). The growth, influenced by genetic and environmental factors, is explained by some non-linear models (Brody, Von Bertalanffy, Gompertz, Logistic, and Richard's) (Kum *et al.* 2010). Briefly, studies on nonlinear growth models provide useful information for breeding intentions (Topal *et al.* 2004; Keskin and Daşkıran 2007; Kucuk *et al.* 2008). Bayram *et al.* (2004) has examined the growth features of Brown Swiss and Friesian cows with only Richard's model. Eyduran *et al.* (2008) reported that Gompertz's model was more effective than Logistic's, monomolecular's and Richard's at early phases of Kivircik and Morkaraman Breeds. Aksoy *et al.* (2011) denoted that the log-linear model was more appropriate than the linear model in Holstein and Brown Swiss Calves. Şahin *et al.* (2014) reported that the growth performance, sexual maturity age, breeding age and appropriate slaughtering age in male and female Anatolian buffalo calves, could be estimated by using Richard's model.

The objective of this study was to examine the live weight of Large White turkeys in both genders, reared in intensive and Free-Ranged systems by some nonlinear models and, to determine the best model.

MATERIAL AND METHOD

This research was carried out in poultry facilities of Agriculture Faculty in Bingol University-Turkey. Totally 60 Large White turkey poults were divided into 2 groups (intensive and Free-with 3 replicates (10 in each).

All the groups were kept inside from 0 to 8 weeks and fed with the feed including 26-28% crude protein and 2800-2900 kcal/kg ME (NRC, 1994). The

birds were fed and watered *adlib*. The groups reared in Intensive system were kept on the ground with litters until the end of 18 weeks fattening period. The lightening program was 23L:1D through the fattening period. Nutritional values and the values of protein and energy of feeds used in the study is shown in Table 1.

Table 1 – Values of protein and energy of feeds used in the study

Feed Type	Week	Protein (%)	ME (kcal/kg)
Starter Feed	0-8	26-28	2800-2900
Grower Feed	9-14	20-23	2900-3000
Finisher Feed	15 - 18	16-19	3000-3200

Free-Range groups had free access to the field in the beginning of the third week, and were held until the end of the fifteenth week of age. A shelter was provided for the birds to keep away from the sunshine during the daytime. Free-Range groups did not receive the concentrated feed during the grazing. The compositions of the grass grazed by the birds were of plants like clover, grass, trefoil, vetch, wheat and barley. After the fifteenth day of fattening, the turkeys were fed the same ration as the indoor groups until the eighteenth weeks of age. The study was carried out between June and October 2014. The birds were wing banded at the beginning of the trail and weighted weekly.

Feeding with intensive and free-range system affected fattening performance of turkeys. Grow growth is also affected by gender. For this reason, the birds were separated by gender and feeding systems and turkeys used in this study were not evaluated all together. This has already been said in the previous sentence.

Von Bertalanffy, Logistic, Gompertz and Gauss models were applied in the weekly analyses of live weights of turkeys in the study.

Von Bertalanffy's growth model is as shown below;

$$Y_t = A * (1 - b * e^{-kt})^3$$

and it has been presented by Von Bertalanffy (1934) with three-parameters, and improved by Beverton & Holt (1957).

Logistic's growth model is expressed as (Graybill & Iyer, 1994);

$$Y_t = A / (1 + e^{-(b+kt)})$$

Gompertz's growth model is expressed as (Winsor, 1932);

$$Y_t = A * e^{-be^{-kt}}$$

Gauss' growth model is expressed as (Norusis, 2005);

$$Y_t = A * (1 - ke^{-bt^2})$$



Here, Y_t is: the weight observed in the age; A is: the asymptotic limit of weight while the age goes to infinity; that is, the highest weight that an animal can reach. This parameter which shows the mature live weight is guessed as the same in all growth curve models; b : the rate of the first weight achieved after birth to mature weight; k is: the speed of getting mature; that is, the speed of the approach of live weight to mature age. The value of $e=2.71828$ is the base of logarithm. Measures such as coefficient of determination (R^2). Adjust R^2 , error mean squares (MSE) and model efficiency, are used in comparing the effectiveness of models impending the observed value

Y_t and expected value $Y_t = E(\hat{Y}_t | X_t)$.

Coefficient of determination (R^2) is;

$$R^2 = \frac{RKT}{GKT} = \frac{\sum_{t=1}^n (\hat{Y}_t - \bar{Y})^2}{\sum_{t=1}^n (Y_t - \bar{Y})^2}$$

It is the total of RKT: Regression mean squares and GKT: General mean squares. The values of Coefficient of determination is between 0-1. As the value approaches to 1, the consistency of the model increases.

Coefficient of determination (Adjust- R^2) is;

$$\bar{R}^2 = 1 - (1 - R^2) \frac{n-1}{n-k}$$

as defined (Gujarati, 2003). Here, n is: number of observation, k : number of parameter.

Error mean squares (MSE) is showed as;

$$MSE = \frac{SSE}{n}$$

SSE: Error sum of squares; n : Observation (Seber & Wild, 1989).

Model efficiency (ME) is expressed as:

$$ME = 1 - \frac{\sum_{t=1}^n (\hat{Y}_t - Y_t)^2}{\sum_{t=1}^n (Y_t - \bar{Y})^2}$$

\hat{Y}_t : Expected value, Y_t : observed value. The efficiency value of the model should be up to 90% for being efficient (Mohanty & Painuli, 2004).

Durbin-Watson statistics (Ünver *et al.* 2011); e_i : Residuals;

$$DW = \frac{\sum_{i=2}^n (e_i - e_{i-1})^2}{\sum_{i=1}^n e_i^2}$$

Akaike Information Criterion (AIC),

$$AIC = n \ln(\hat{\sigma}^2) + 2M$$

Schwartz Bayesian Criterion (SBC);

$$SBC = n \ln(\hat{\sigma}^2) + M \ln(n)$$

as defined (Wei, 2006). Here, $\hat{\sigma}^2$ is the maximum likelihood of σ , M is the number of parameters in the model, and n is the effective number of observations.

A , B and k parameters, were determined as a result of Levenberg Marquardt iteration technique by using SPSS 22.0 statistical program. While iteration was done, $1.0E-8$ was used as convergence criteria (Akbaş *et al.* 2001).

In comparing the models, the measures of coefficient of determination (R^2), Error mean squares (MSE) which shows the difference between the points belongs to true growth and growth curve determined to the model, and model efficiency were used.

RESULTS AND DISCUSSION

Live weight averages of male and female turkeys in the weeks of 0-18 is shown in Table 2.

Table 2 – Live weights (g) of male and female turkey's growth in Intensive and free-range systems according to the weeks

Age (Weeks)	Intensive System		Free-Range System	
	Body Weight (g)-Male	Body Weight (g)-Female	Body Weigh (g)-Male	Body (g)- Female
0	63.10	61.87	64.75	65.14
2	234.71	224.75	262.18	249.82
4	687.14	642.40	754.64	689.06
6	1685.00	1509.25	1768.27	1517.06
8	3188.14	2854.05	3460.73	2928.24
10	5249.00	4597.30	3448.73	2933.82
12	6920.57	5945.60	4405.73	3757.94
14	8186.43	7369.30	5308.55	4491.71
16	10833.57	9080.00	7650.55	6401.24
18	12285.71	10164.20	8438.18	6985.29



Hatching weight for male and female birds were 63.10 and 61.87g in commercial groups, 64.75 and 65.14 g in Free-Range groups, respectively. While the live weights of the turkeys for male and female were 12285.71 and 10164.20 g in commercial groups, and 8438.18 g and 6985.29 g in Free-Range groups at the end of the trail, respectively.

Live weight values of turkeys were comparatively observed by Von Bertalanffy, Logistic, Gompertz and Gauss models, and the results are shown in Table 3.

As it is seen in Table 3, the turkeys raised in intensive system had the values of Error mean squares (MSE) = 67085.950, Coefficient of correlation (r) = 0.9994, and growth speed = 0.166 g according to Von Bertalanffy's model. Determination coefficients (R^2) = 0.9974, Adjusted- R^2 = 0.9967, Model efficiency (ME) = 0.9974, DW = 2.335, AIC = 136.596 and SBC = 143.656. According to the results obtained with the Logistic's model were MSE = 178332.669, r = 0.997 and growth speed 0.610 g. R^2 = 0.996, Adj- R^2 = 0.994, ME = 0.990, DW = 1.491, AIC = 146.373 and SBC = 154.410. The results obtained with Gompertz's model were MSE = 79876.142, r = 0.998 and growth speed 0.279 g. R^2 = 0.9969, Adj- R^2 = 0.9960, and ME = 0.9969, DW = 2.416, AIC = 138.341 and SBC = 145.576. The results obtained with Gauss's model were MSE = 74646.109, r = 0.999 and growth speed 0.011 g, R^2 = 0.9971, Adj- R^2 = 0.9962, and ME = 0.9971, DW = 2.587, AIC = 137.664 and SBC = 144.831.

In female turkeys grown in intensive system; in Von Bertalanffy's model, the results were calculated as MSE = 15477.841, r = 0.99960, the growth speed = 0.200 g. R^2 = 0.99915, Adjusted- R^2 = 0.99891, model efficiency (ME) = 0.99940, DW = 2.441, AIC = 121.931 and SBC = 127.524. In Logistic's model, the results were calculated as MSE = 71695.646, r = 0.9983, and the growth speed = 0.648 g. R^2 = 0.9961, Adj- R^2 = 0.9949, and ME = 0.9972, DW = 1.141, AIC = 137.261 and SBC = 144.387. In Gompertz's model, the results were calculated as MSE = 16176.236, r = 0.99956, and the growth speed = 0.313 g. R^2 = 0.99911, Adj- R^2 = 0.99886, and ME = 0.99937, DW = 2.280, AIC = 122.372 and SBC = 128.009. In Gauss's model, the results were calculated as MSE = 24691.170, r = 0.99932, and the growth speed = 0.014 g. R^2 = 0.9986, Adj- R^2 = 0.9983, and ME = 0.99904, DW = 1.864, AIC = 126.601 and SBC = 132.661.

According to the Gompertz's model, growth speeds in different studies were 0.2137 in male partridges and 0.2255 in female partridges (Çetin et al., 2007); 0.010 in sheep (Tariq et al., 2013), 0.218 in male calves and 0.309 in female calves (Şahin et al. (2014), 0.010 in female Anatolian buffalo and 0.007 in male Anatolian buffalo (Sosyal et al. 2015). According to the Von Bertalanffy's model, the growth speeds were 0.0044 in sheep (Tariq et al., 2013) and 0.005 in male buffalo and 0.012 in female buffalo (Soysal et al. 2015).

In Von Bertalanffy's model, male turkeys reared in Free-Range system the results were found as follow; MSE = 231089.161, r = 0.9897 and growth speed = 0.093, R^2 = 0.9794, adjusted- R^2 = 0.9735, model efficiency (ME) = 0.9794, DW = 2.089, AIC = 148.965 and SBC = 157.261. In the results obtained in Logistic's model, MSE = 313900.941, r = 0.9897, growth speed = 0.456 g, R^2 = 0.972, Adj- R^2 = 0.964, ME = 0.972, DW = 1.616, AIC = 152.027 and SBC = 160.630. According to Gompertz's model, the results were; MSE = 254176.033, r = 0.9888, and the growth speed = 0.185g, R^2 = 0.9773, Adj- R^2 = 0.9709, ME = 0.9773, DW = 1.932, AIC = 149.917 and SBC = 158.309. In Gauss's model, the results were; MSE = 267465.980, r = 0.988, and the growth speed is found as 0.010 g. R^2 = 0.9761, Adj- R^2 = 0.9693, ME = 0.9761, DW = 1.846, AIC = 150.427 and SBC = 158.869.

In Von Bertalanffy's model for female turkeys reared in Free-Range system, the results were; MSE is 151000.416, r = 0.9902, and the growth speed = 0.102g, R^2 = 0.9804, Adjusted- R^2 = 0.9748, model efficiency (ME) = 0.9802, DW = 2.167, AIC = 144.709 and SBC = 152.580. In Logistic's model; MSE = 208664.111, r = 0.9868, and growth speed = 0.461, R^2 = 0.9729, Adj- R^2 = 0.9651, ME = 0.9727, DW = 1.655, AIC = 147.944 and SBC = 156.138. In Gompertz's model; MSE = 166554.915, r = 0.9893, and growth speed = 0.194 g, R^2 = 0.9784, Adj- R^2 = 0.9722, ME = 0.9782, DW = 1.998, AIC = 145.690 and SBC = 153.659. In Gauss's model; MSE = 177604.528, r = 0.9884, growth speed = 0.012 g, R^2 = 0.977, Adj- R^2 = 0.9704, ME = 0.9768, DW = 1.888, AIC = 146.332 and SBC = 154.365. In all models, the most appropriate model, in which MSE value is the least, was selected.

Mathematical growth models of male and female turkeys reared in intensive and Free-Range systems is shown in Table 4.



Table 3 – Parameters of growth models for male and female Large White Turkeys

Intensive system-Male

Parameters	Von Bertalanffy	Logistic	Gompertz	Gauss
A	25673.654	14233.320	19131.256	21597.618
b	0.968	-3.756	5.490	0.011
k	0.166	0.610	0.279	1.003
r	0.9994	0.997	0.998	0.999
R ²	0.9974	0.9931	0.9969	0.9971
Adj R ²	0.9967	0.9911	0.9960	0.9962
MSE	67085.950	178332.669	79876.142	74646.109
ME	0.9974	0.9931	0.9969	0.9971
DW	2.355	1.491	2.416	2.587
AIC	136.596	146.373	138.341	137.664
SBC	143.656	154.410	145.576	144.831

Intensive system-Female

Parameters	Von Bertalanffy	Logistic	Gompertz	Gauss
A	17583.723	11246.675	14213.138	15308.655
b	1.003	-3.752	5.629	0.014
k	0.200	0.648	0.313	1.005
r	0.99960	0.9983	0.99956	0.99932
R ²	0.99915	0.9961	0.99911	0.9986
Adj R ²	0.99891	0.9949	0.99886	0.9983
MSE	15477.841	71695.646	16176.236	24691.170
ME	0.99940	0.9972	0.99937	0.99904
DW	2.441	1.141	2.28	1.864
AIC	121.931	137.261	122.372	126.601
SBC	127.524	144.387	128.009	132.661

Free-Range system- Male

Parameters	Von Bertalanffy	Logistic	Gompertz	Gauss
A	21771.690	11769.879	18625.261	14594.647
b	0.823	-3.171	4.194	0.010
k	0.093	0.456	0.185	0.979
r	0.9897	0.9864	0.9888	0.988
R ²	0.9794	0.972	0.9773	0.9761
Adj R ²	0.9735	0.964	0.9709	0.9693
MSE	231089.161	313900.941	254176.033	267465.980
ME	0.9794	0.972	0.9773	0.9761
DW	2.089	1.616	1.932	1.846
AIC	148.965	152.027	149.917	150.427
SBC	157.261	160.630	158.309	158.869

Free-Range system- Female

Parameters	Von Bertalanffy	Logistic	Gompertz	Gauss
A	16614.034	9435.685	14263.744	11175.027
b	0.812	-3.104	4.089	0.012
k	0.102	0.461	0.194	0.976
r	0.9902	0.9868	0.9893	0.9884
R ²	0.9804	0.9729	0.9784	0.9770
Adj R ²	0.9748	0.9651	0.9722	0.9704
MSE	151000.416	208664.111	166554.915	177604.528
ME	0.9802	0.9727	0.9782	0.9768
DW	2.167	1.655	1.998	1.888
AIC	144.709	147.944	145.690	146.332
SBC	152.580	156.138	153.659	154.365

A, b, k: Model parameters, (A: Asymptotic weight; b: Integration constant, k: Maturing index, r: Correlation coefficients, R²: Determination coefficients, Adj R²: Adjusted determination coefficients, MSE: Mean of Squares Error, ME: Model efficiency, DW: Durbin-Watson, AIC: Akaike Information Criterion, SBC: Schwartz Bayesian Criterion (SBC).



Table 4 – Functions belonging to growth curves of turkeys

Intensive system			
Models	Functions		
	Male		Female
Von Bertalanffy	$Y = 25673.654(1 - 0.968e^{-0.166t})^3$		$Y = 17583.723(1 - 1.003e^{-0.200t})^3$
Logistic	$Y = 14233.320/(1 + e^{-3.756+0.610t})$		$Y = 11246.675/(1 + e^{-3.752+0.648t})$
Gompertz	$Y = 19131.256(e^{-5.490e^{-0.279t}})$		$Y = 14213.138(e^{-5.629e^{-0.313t}})$
Gauss	$Y = 21597.618(1 - 1.003e^{-0.011t^2})$		$Y = 15308.655(1 - 1.005e^{-0.014t^2})$
Free-Range system			
Model	Function		
	Male		Female
Von Bertalanffy	$Y = 21771.690(1 - 823e^{-0.093t})^3$		$Y = 16614.034(1 - 812e^{-0.102t})^3$
Logistic	$Y = 11769.879/(1 + e^{-3.171+0.456t})$		$Y = 9435.685/(1 + e^{-3.104+0.461t})$
Gompertz	$Y = 18625.261(e^{-4.194e^{-0.185t}})$		$Y = 14263.744(e^{-4.089e^{-0.194t}})$
Gauss	$Y = 14594.647(1 - 0.979e^{-0.010t^2})$		$Y = 11175.027(1 - 0.976e^{-0.012t^2})$

When we compare growth models of male turkeys reared in Intensive system, the values for Von Bertalanffy's, Logistic's, Gompertz's, and Gauss's models were; MSE=67085.950, 178332.669, 79876.142, and 74646.109; coefficient of determinations (R^2)=0.9974, 0.9931, 0.9969, and 0.9971; adjusted coefficient of determinations (Adjust- R^2)= 0.9967, 0.9911, 0.9960, and 0.9962, respectively. Again for the same models and in the same order; ME were 0.9967, 0.9911, 0.9960, and 0.9962. DW were 2.355, 1.491, 2.416, and 2.587; AIC were 136.596, 146.373, 138.341, and 137.664; SBC were 143.656, 154.410, 145.576, and 144.831, respectively. Von Bertalanffy's model had the lowest MSE AIC and SBC values, and highest R^2 , Adj- R^2 , and ME values; and Logistic's model was the one which had the highest MSE, AIC and SBC values and the lowest R^2 , Adj- R^2 , and ME values. Thus, the most relevant model for estimation of growth curve was Von Bertalanffy's growth model. In other words, it was determined that Von Bertalanffy's growth model identifies the growth curve of male turkeys in intensive breeding system best, and Logistic's model identifies the growth curve the least.

Comparing growth models for female turkeys reared in intensive system, the values for Von Bertalanffy's, Logistic's, Gompertz's, and Gauss' models were; MSE=15477.841, 71695.646, 16176.236, and 24691.170; coefficient of determinations (R^2)= 0.99915, 0.9961, 0.99911, and 0.9986; adjusted

coefficient of determinations (Adjust- R^2)= 0.99891, 0.9949, 0.99886, and 0.9983, respectively. Again for the same models in same order; ME= 0.9994, 0.9972, 0.9993, and 0.99904. DW were 2.441, 1.141, 2.280 and 1.864; AIC were 121.931, 137.261, 122.372 and 126.601; SBC were 127.524, 144.387, 128.009 and 132.661, respectively. Von Bertalanffy's model had the lowest MSE, AIC and SBC values and highest R^2 , Adj- R^2 , and ME values; and Logistic's model was the one which had the highest MSE, AIC and SBC values and the lowest R^2 , Adj- R^2 , and ME values. Thus, the most relevant model for estimation of growth curve of female turkeys was seen in Von Bertalanffy's growth model. In other words, the best model was Von Bertalanffy's model and the worst one was Logistic's model for female turkeys in intensive breeding system. The results obtained in coefficient of determinations in accordance with the results of approximate (0.99) coefficient of determinations were Gompertz's, Logistic's, Richard's and Morgan-Mercer-Flodin's models in growth curves of Large White turkeys.

When we compare growth models for male turkeys reared in Free-Range system, the calculated values for Von Bertalanffy's, Logistic's, Gompertz's, and Gauss' models were; MSE=231089.161, 313900.941, 254176.033, and 267465.980; coefficient of determinations (R^2)= 0.9794, 0.972, 0.9773, and 0.9761; adjusted coefficient of determinations



(Adjust-R²)= 0.9735, 0.964, 0.9709, and 0.9693, respectively. Again for the same models in same order, ME were 0.9794, 0.972, 0.9773, and 0.9761. DW were 2.089, 1.616, 1.932 and 1.846; AIC were 148.965, 152.027, 149.917 and 150.427; SBC were 157.261, 160.630, 158.309 and 158.869, respectively. Von Bertalanffy's model had the lowest MSE, AIC and SBC values and highest R², Adj-R², and ME values; however, Logistic's model had the highest MSE, AIC and SBC values and the lowest R², Adj-R², and ME values. In other words, Von Bertalanffy's growth model best identifies growth curve values and Logistic's model identifies the growth curves the least for male turkeys reared in Free-Range system.

When we compare growth models for female turkeys, the calculated values for Von Bertalanffy, Logistic, Gompertz, and Gauss models were: MSE=151000.416, 208664.111, 166554.915, and 177604.528; coefficient of determinations (R²) = 0.9804, 0.9729, 0.9784, and 0.9770; adjusted coefficient of determinations (Adjust-R²) = 0.9748, 0.9651, 0.9722, and 0.9704, respectively. For the same models in the same order, ME= 0.9802, 0.9727, 0.9782, and 0.9768. DW were 2.167, 1.655, 1.998 and 1.888; AIC were 144.709, 147.944, 145.690 and 146.332; SBC were 152.580, 156.138, 153.659 and 154.365, respectively. Von Bertalanffy model had the lowest MSE, AIC and SBC values and highest R², Adj-R², and ME values; but Logistic model had the highest MSE, AIC and SBC values and the lowest R²,

Adj-R², and ME values. Thus, the most relevant model for estimation of growth curve of live weight of female turkeys was Von Bertalanffy growth model, and the least relevant was Logistic model. Similar results for Von Bertalanffy model were also reported by Akbaş *et al.* (2001) in Holstein Friesians cows, and by Soysal *et al.* (2015) in male buffalo. However, the best fitted model in Anatolian buffalo was Richards model (Şahin *et al.* 2014).

The results obtained in coefficient of determinations were 0.99 with Gompertz, Logistic, Richards, Morgan-Mercer-Flodin models in growth curves of Large White turkeys. In both of the systems, Von Bertalanffy model had the highest mature weight, and, Logistic model had the least in both gender.

In this study, estimation curves identified with 4 different growth models were in Figure 1 and 2 for male and female turkeys reared in Intensive system, and in Figure 3 and 4 for male and female turkeys reared in Free-Range system, respectively. Even if the values observed in the models were so close to each other, Von Bertalanffy model was more relevant than the others.

Many measures such as the consistence of growth models, R², MSE, their biologic relevancies, and consistency of estimations can generally be used in comparison of growth curve models characterizing the growth in livestock production (Torres and Ortiz, 2005; Malhado *et al.*, 2008, Torres *et al.*, 2012).




Figure 1 – Growth curves of male Large White turkeys in Intensive system




Figure 2 – Growth curves of female Large White turkeys in Intensive system




Figure 3 – Growth curves of male Large White turkeys in Free-Range system

In the comparison of growth curve models in which R^2 and MSE values were scored together, the model which had the highest R^2 value and the least MSE values, explain the live weight changes according to the age. (Quirino et. al. 1999; Bayram % Akbulut; 2009; Yıldız et al. 2009; Kucuk & Eyduran 2009; Mohammed, 2015).

Çetin et al. (2007), calculated DW values for Gompertz, Richards and Logistic as 1.9543, 1.6179, 0.4017 in male partridges; 1.1763, 1.2025 and 0.5426 in female partridges, respectively. Faridi et al. (2011), AIC were 5980.53, 5632.6, 5525.255, 5912.12 and 5713.70; SBC were 6002.34, 5658.82, 5555.45, 5920.85 and 5726.80 by using NT1, NT2, NT3, Gompertz




Figure 4 – Growth curves of female Large White turkeys in Free-Range system

and Richards models for growth in broiler breeder flocks, respectively. Narinc *et al.* (2013) calculated AIC as -301.43, -299.56, -282.51 and -307.47; SBC as -295.57, -291.75, -274.71 and -297.71 for Gamma, McNally, Modified compartmental and Adams-Bell models in egg production in Japanese quail, respectively. Besides AIC were found as -176.75, -195.92, -356.32 and -386.56; SBC were as -171.02, -188.27, -346.76 and -378.95, respectively by using Gamma, McNally, Modified compartmental and Adams-Bell models in egg production (Narinc *et al.* (2014) .

CONCLUSION

In this research, growth models of male and female turkeys reared in Intensive and Free-Range systems from their hatching to the 18th week of age were compared. The 4 models have been applied for live weight data of turkeys. The values of R^2 and $Adj-R^2$ of all the models have been found very high. The efficiency value of the model was up to 0.90 in all models and that means that models were efficient.

In conclusion, Von Bertalanffy model explained the weight-age change of male and female turkeys reared in different systems best.

REFERENCES

Abegaz S, Van Wyk JB, Olivier JJ. Estimation of genetic and phenotypic parameters of growth curve and their relationship with early growth and productivity in Horro sheep. *Archiv Tierzucht* 2010;53(1):85–94.

Aggrey SE. Comparison of three nonlinear and spline regression models for describing chicken growth curves. *Poultry Science* 2002;81(12):1782–1788.

Agudelo-Gómez D, Hurtado-Lugo N, Cerón-Muñoz MF. Growth curves and genetic parameters in colombian buffaloes (*Bubalus bubalis* Artiodactyla, Bovidae). *Revista Colombiana Ciencias Pecuarias* 2009;22(2):178-188.

Akbaş Y, Taşkın T, Demirören E. Comparison of several models to fit the growth curves of kıvrıkcık and dağlıç male lambs. *Turkish Journal of Veterinary and Animal Science* 1999;23(Suppl 3):537-544.

Akbaş Y, Akbulut Ö, Tüzemen N. Growth of Holstein cows in high altitude of Turkey. *Indian Journal Animal Science* 2001;71(5):476-479.

Aksoy A, Külekçi M, Bayram B, Akbulut Ö. Linear and log-linear model approach on determining the factors affecting live weight gain of holstein and brown swiss calves. *Journal of Agricultural Faculty of Atatürk University* 2011;41(2):123-127.

Alpan O, Arpacık R. Sığır yetiştiriciliği. Baskı: Şahin Matbaası; 1998.

Akçapınar H, Özbeyaz C. Hayvan Yetiştiriciliği Temel ilkeleri. Ankara: Kariyer Matbaacılık; 1999.

Ayied AY, Al-Hello F, Al-Jassim EF. Growth patterns of Arabi sheep in South of Iraq. *Journal of Tikrit University for Agricultural Science* 2011;11:384–391

Bayram B, Akbulut Ö. Analysis of growth curve in brown swiss and holstein friesian cattle using linear and non-linear models. *Hay Üretim* 2009;50(2):33-40.

Bayram B, Akbulut Ö, Yanar M, Tüzemen N. Analysis of growth characteristics using the richards model in female brown swiss and holstein friesian cattle. *Turkish Journal of Veterinary and Animal Scienca* 2004;28: 201-208.

Behr V, Hornick JL, Cabaraux JF, Alvarez A, Istasse L. Growth patterns of Belgian Blue replacement heifers and growing males in commercial farms. *Livestock Production Science* 2001;71(2-3):121-130.



- Bertalanffy LV. Wandlungen des biologischen Denkens, Neue Jahrbücher für Wissenschaft und Jugendbildung 1934;10:339-366.
- Beverton RJH, Holt SJ. On the dynamics of exploited fish populations. London : Her Majesty's Stationery Office, 1957; v.19, p.533.
- Bilgin OC, Emsen E, Davis ME. Comparison of nonlinear models for describing growth of scrotal circumference in Awassi male lambs. Small Ruminant Research 2004;52:155-160.
- Çetin M, Şengül T, Söğüt B, Yurtseven S. Comparison of growth models of male and female partridges. Journal of Biological Sciences 2007;7:964-968.
- Doğan, İ. The Evaluation of Growth of Lambs by Multidimensional Scaling. Uludağ Üniversitesi Journal of the Faculty Veterinary Medicine 2003;22(1-3):33-37.
- Eleroglu H, Yıldırım A, Şekeroğlu A, Çoksöyler FN, Duman M. Comparison of growth curves by growth models in slow-growing chicken genotypes raised the organic system. International Journal of Agriculture and Biology 2014;16:529-535.
- Enders, W. Applied econometric time series. 3rd ed. New York: John Wiley & Sons; 2010.
- Eyduran E, Kucuk M, Karakus K, Ozdemir T. New approaches to determination of the best nonlinear describing growth at early phases of kivircik and morkaraman breeds. Journal of Animal and Veterinary Advances 2008;7(7):799-804.
- Faridi A, Mottaghtalab, Rezaee F, France J. Narushin-Takma models as flexible alternatives for describing economic traits in broiler breeder flocks. Poultry Science 2011;90:507-515.
- Graybill, FA, Iyer, H K. Regression analysis: concepts and applications. New York: Hardcover; 1994.
- Gujarati, DN. Basic econometrics. 4 th ed. New York: McGraw-Hill; 2003.
- Kucuk M., Eyduran E. The determination of the best growth model for akkaraman and german lackheaded mutton x akkaraman b₁ crossbred lambs. Bulgarian Journal of Agricultural Science 2009;15(1):90-92.
- Kum D, Karakus K, Ozdemir T. The best nonlinear function for body weight at early phase of Norduz female lambs. Trakia Journal Science 2010;8(2):62-67.
- Lambe NR, Navajas EA, Simm G, Bünger LA genetic investigation of various growth models to describe growth of lambs of two contrasting breeds. Journal of Animal Science 2006;84:2642-2654.
- Malhado, CHM, Ramos AA, Carneiro PLS, Souza JC, Wechsler FS, Eler JP, et al. Nonlinear models to describe the growth of the buffaloes of murrh breed. Archives Zootecnia 2008;57(220):497-503.
- Mohammed, F. A. Comparison of three nonlinear functions for describing chicken growth curves. Scientific Agriculture 2015;9(3):120-123.
- Mohanty M., Painuli, D. K. Modelling rice seedling emergence growth and under tillage and residue management in a rice-wheat system on vertisol in Central India. Soil and Tillage Research 2004;76:167-174.
- Narinc D, Karaman E, Aksoy T, Firat MZ. Investigation of nonlinear models to describe long-term egg production in Japanese quail. Poultry Science 2013;92:1376-1682.
- Narinc D, Uckardes F, Aslan E. Egg production curve analyses in poultry science. World's Poultry Science Journal 2014;70:817-828.
- Ngeno K, Bebe BO, Kahi AK. Estimation of growth parameters of indigenous chicken populations intensively reared in Kenya. Egerton Journal of Science Technology 2010;11:13-28
- Norris D, Ngambi JW, Benyi K, Makgah-lela, ML, Shimelis HA, Nesamvuni, EA. Analysis of growth curves of indigenous male venda and naked neck chickens. South African Journal of Animal Science 2007;37:21-26.
- Norusis MJ. SPSS 14.0 advanced statistical procedures companion. New Jersey: Prentice Hall; 2005.
- NRC. Nutrient Requirements of Poultry. 9th ed. Washington: National Academy Press; 2005.
- Prestes AM, Garnero ADV, Marcondes CR, Damé MC, Janner EA, Rorato PRN. Estudo da curva de crescimento de bubalinos da raça Murrh criados no estado do Rio Grande do Sul. Anais do 9º Simpósio Brasileiro de Melhoramento Animal; 2012 Jun 20-26; João Pessoa: ABMA; 2012.
- Quirino CR, Bergmann JA, Vale Filho, VR, Andrade VJ, Pereira, JC. Evaluation of four mathematical functions to describe scrotal circumference maturation in Nelore bulls. Theriogenology 1999;52(1):25-34.
- Reddish JM, Lilburn MS. A comparison of growth and development patterns in diverse genotypes of broilers. Pullet growth. Poultry Science 2004;83:1072-1076.
- Rizzi C, Contiero B, Casandro M. Growth patterns of Italian local chicken populations. Poultry Science 2013;92(8):2226-2235.
- Saghi DA, Aslaminejad A, Tahmoospur M, Farhangfar H, Nassiri M, Dashab GR. Estimation of genetic parameters for growth traits in Baluchi sheep using Gompertz growth curve function. Indian Journal of Animal Science 2012;82:889-892.
- Schulze V, Rohe R, Looft H, Kalm E. Genetic analysis of the course of individual growth and feed intake of group-penned performance tested boars. Archiv Tierzucht 2001;44:139-156.
- Seber GAF, Wild CJ. Nonlinear regression. New York: John Wiley and Sons; 1989.
- Soysal MI, Gurcan EK, Genc S, Aksel M. The Comparison of growth curve with different models in anatolian buffalo. Journal of Tekirdag Agricultural Faculty 2015;12(3):57-61.
- Şahin A, Ulutaş Z, Karadavut U, Yıldırım A, Arslan S. Comparison of growth curve using some nonlinear models in anatolian buffaloe calves. Kafkas Üniversitesi Veteriner Fakültesi Dergisi 2014;20(3):357-362..
- Şengül T, Kiraz S. Ağır Beyaz hindilerde büyüme eğrilerinin tanımlanmasında doğrusal olmayan modeller. Turkish Journal of Veterinary and Animal Science 2005;29:331-337.
- Tariq MM, Bajwa MA, Waheed A, Eyduran E, Abbas F, Bokhari FA, et al. Growth curve in Mengali sheep breed of Bachistan. Journal of Animal and Plant Science 2011;21:5-7.
- Tariq, MM, Iqbal F, Eyduran E, Bajwa MA, Huma ZE, Waheed A. Comparison of Non-Linear Functions to Describe the Growth in Mengali Sheep Breed of Balochistan. Pakistan Journal of Zoology 2013;45(3):661-665.
- Topal M, Ozdemir M, Aksakal V, Yıldız N, Dogru U. Determination of the best nonlinear function in order to estimate growth in Morkaraman and Awassi lambs. Small Ruminant Research 2004;55:229-232.
- Torres V, Ortiz J. Application of modelling and simulation to the production and feeding farm animals. Cuban Journal of Agricultural Science 2005;39:385.
- Torres V, Barbosa I, Meyer R, Noda A, Sarduy L. Criteria of goodness of fit test in the selection of non-linear models for the description of biological performances, Cuban Journal of Agricultural Science 2012;46(4):345-350.
- Ünver Ö, Gamgam H, Altunkaynak B. SPSS uygulamalı temel istatistik yöntemler. Seçkin: Yayıncılık San. Ve Tic. A.Ş; 2011.
- Winsor CP. The Gompertz curve as a growth curve. Proceedings of the National Academy of Science 1932;18(1):1-8
- Yıldız G, Soysal M İ, Gürçan EK. Determination of growth curve with different model in the Karacabey Merinos xKivircik crossbred lambs raised in Tekirdağ provinces. Tekirdağ Ziraat Fakültesi Dergisi 2009;6(1):11-19.