

Factors associated to hypoxemia in patients undergoing coronary artery bypass grafting

Fatores associados à ocorrência de hipoxemia em pacientes submetidos à revascularização miocárdica

Natália Pereira dos Santos¹, PT; Rodrigo Martins Mitsunaga¹; Daniel Lago Borges¹, PT, MsC; Marina de Albuquerque Gonçalves Costa¹, PT; Thiago Eduardo Pereira Baldez¹, PT; Ilka Mendes Lima¹, PT; Karol Cristina Fonseca Moura¹, PT; Josimary Lima da Silva Lula²

DOI: 10.5935/1678-9741.20130056

RBCCV 44205-1482

Abstract

Introduction: Hypoxemia is a frequent pulmonary complication in the postoperative coronary artery bypass graft. Detection of factors associated with their occurrence may indicate patients at risk for this complication, which allows tracing specific therapeutic and consequently reduce morbidity and mortality.

Objective: To identify related factors to hypoxemia occurrence in immediate coronary artery bypass graft postoperative.

Methods: In this retrospective cohort study, we studied 100 patients submitted to elective on-pump artery bypass graft, between April 2010 and December 2011, at a reference university hospital for cardiac surgery in the state of Maranhão. It was considered hypoxemia gas exchange ratio less than or equal to 300 mmHg. Associated variables with perioperative hypoxemia were defined by the Student T test, G or Mann-Whitney tests, Chi-square, or Fisher's exact test and multiple linear regression.

Results: Among studied variables, high body mass index ($P=0.036$) and smoking ($P=0.024$) were significantly associated with hypoxemia in the immediate coronary artery bypass graft postoperative. Hypoxemia incidence in this period was 55% and

did not affect mechanical ventilation duration and Intensive Care Unit length of stay.

Conclusion: In this sample, body mass index and smoking were associated to hypoxemia. These data reinforce the importance of clinical assessment to identify patients at risk for this complication, considering its high incidence in immediate postoperative period.

Descriptors: Oxygen level. Postoperative period. Myocardial revascularization.

Resumo

Introdução: A hipoxemia é uma complicação pulmonar frequente no pós-operatório de revascularização miocárdica. A detecção de fatores associados a sua ocorrência pode indicar pacientes de risco para essa complicação, o que possibilita traçar terapêuticas específicas e, conseqüentemente, diminuir a morbimortalidade.

Objetivo: Identificar fatores relacionados à ocorrência de hipoxemia no pós-operatório imediato de revascularização miocárdica.

Métodos: Nesta coorte retrospectiva, foram estudados 100

¹ Universidade Federal do Maranhão, Hospital Universitário, São Luís, MA, Brazil.

² Hospital São Domingos, São Luís, MA, Brazil.

Correspondence address:

Daniel Lago Borges

Rua Barão de Itapary, 227 – Centro – São Luís, MA, Brazil

Zip code: 65020-070

E-mail: dlagofisio83@hotmail.com

Abbreviations, acronyms & symbols	
AMI	Acute myocardial infarction
ARF	Acute renal failure
BMI	Body mass index
CABG	Coronary artery bypass graft
CAD	Coronary atherosclerotic disease
COPD	Chronic obstructive pulmonary disease
CPB	Cardiopulmonary bypass
CTA	Previous angioplasty
DLP	Dyslipidemia
DM	Diabetes mellitus
FiO ₂	Inspired oxygen fraction
ICU	Intensive care unit
IPO	Immediate postoperative
LITA	Left internal thoracic artery
LV	Left ventricular
PaO ₂	Pressure of oxygen in the arterial blood
PEEP	Positive end-expiratory pressure
RR	Respiratory rate
VT	Volume tidal
WHO	World Health Organization

pacientes que se submeteram à cirurgia eletiva de revascularização miocárdica com uso de circulação extracorpórea, no período de abril de 2010 a dezembro de 2011, no Hospital Uni-

INTRODUCTION

Coronary artery bypass grafting (CABG) surgery, despite all the advances in clinical treatment and percutaneous interventions, is still used in coronary heart disease treatment [1]. Pulmonary complications in cardiac surgery postoperative period are very frequent and represent significant morbidity and mortality cause [2].

Hypoxemia is one of the main causes of pulmonary complications characterized by decrease of arterial oxygen partial pressure (PaO₂) [3,4]. Measuring severity of the pulmonary injury in mechanically ventilated patients is performed by PaO₂ and inspired oxygen fraction (FiO₂) ratio [5].

Diverse mechanisms have been accounted for its development on cardiac surgery postoperative, especially atelectasis, shunt increase, respiratory and thoracic mechanics alterations, capillary and pulmonary parenchyma changes secondary to the left ventricular (LV) dysfunction or pulmonary endothelium injury [3,6].

Elderly, obese and patients with left ventricular dysfunction (ejection fraction of the left ventricle < 55%) and prolonged cardiopulmonary bypass (CPB) time (> 120 minutes) had higher risks to postoperative surgery hypoxemia [7]. Pulmonary chronic diseases, diabetes and preoperative acute myocardial infarction are independent risk factors for hypoxemia after CABG [8].

versitário de referência para cirurgia cardíaca no estado do Maranhão. Considerou-se hipoxemia índice de troca gasosa menor ou igual a 300 mmHg. A associação das variáveis perioperatórias com a ocorrência de hipoxemia foi definida por meio dos testes t de Student ou Mann Whitney, Qui-quadrado, teste G, ou Teste de Fisher, além de regressão linear múltipla.

Resultados: Dentre as variáveis estudadas, o índice de massa corpórea elevado ($P=0,036$) e o tabagismo ($P=0,024$) apresentaram associação estatisticamente significativa com a ocorrência de hipoxemia no pós-operatório imediato de revascularização miocárdica. A incidência de hipoxemia no pós-operatório imediato foi 55%. A ocorrência de hipoxemia não interferiu nos tempos de ventilação mecânica e de estadia na Unidade de Terapia Intensiva.

Conclusão: Nesta amostra, houve associação entre o índice de massa corpórea e o tabagismo com a ocorrência de hipoxemia. Esses dados reforçam a importância da avaliação clínica para identificação do paciente de risco para ocorrência dessa complicação, já que esta apresenta elevada incidência no pós-operatório imediato.

Descritores: Nível de oxigênio. Período pós-operatório. Revascularização miocárdica.

According to a recent study [7], predictor factors detection may indicate patients at high risk for postoperative hypoxemia that demand mechanical ventilation strategies, in order to prevent and treat pulmonary collapse during intra-operative period, to minimize hypoxemia impact on mechanical ventilation duration, to reduce morbidity and mortality related to cardiac intervention, to reduce length of stay at Intensive Care Unit (ICU), and hospital costs.

This study is designed to analyze factors related to hypoxemia in a specific population of patients undergoing CABG surgery, to determine its incidence in immediate post-operative period, and to evaluate if hypoxemia interfered on mechanical ventilation duration and length of stay in ICU.

METHODS

It is a retrospective descriptive cohort study conducted in a reference university hospital for cardiac surgery in São Luis, MA, Brazil, after Research Ethics Committee approval (Protocol N° 0051/2012), as required by Resolution N° 196/96 of the National Health Council.

Patients who underwent CABG between April 2010 and December 2011 and were admitted to the Cardio ICU in the referred hospital were included in research.

Adult patients (> 18 years old) from both genders were included in the study, with no prior cardiac surgery,

submitted to elective on-pump CABG not associated to other procedures and ventilated in assisted/controlled volume mode, with PEEP (positive end-expiratory pressure) 8 cmH₂O and FiO₂ 40%.

Missing data related to PaO₂/FiO₂ ratio, perioperative death and returning to operation room were exclusion criteria to this study.

Data were collected from Physical Therapy Evaluation form – divided in pre, intra and postoperative periods, filled with information from medical records – and registered in a specifically form designed for this study.

During study period, two hundred and fifty-five patients were admitted to Cardiovascular ICU after CABG (Figure 1). Among them, 100 patients fulfilled all criteria and general characteristics are described in Table 1.

Preoperative data

Preoperative variables were: age, considering as advanced age ≥ 60 years old; gender; nutritional status, determined by Body Mass Index (BMI) calculated by weight/height² ratio, considered as eutrophic = 20 to 24.9 kg/m², overweight = 25 to 29.9 kg/m² and obese ≥ 30 kg/m², established by the World Health Organization (WHO) [9].

Other studied variables were risk factors to coronary diseases registered in anesthesiologist assessment available in medical records: diabetes mellitus (DM), hypertension, dyslipidemia (DLP), smoking (also were considered former smokers) and associative diseases: chronic obstructive pulmonary disease (COPD), acute renal failure (ARF), acute myocardial infarction (AMI) and prior angioplasty.

It was not possible to correlate left ventricular ejection fraction (left ventricular dysfunction evaluation) due lack of data, cause it is not a routine service to have an echocardiogram in all preoperative patients.


Intraoperative data

For CABG patients underwent general anesthesia, tracheal intubation, controlled mechanical ventilation, surgical access via median sternotomy and CPB, performed with graphs of the left internal thoracic artery (LITA) and/or saphenous vein and the placement of mediastinal and pleural drains.

Intraoperative analyzed variables were numbers of graphs, LITA use, number of drains, surgery time, considering a prolonged intervention ≥ 210 minutes; pump time, considering prolonged when above 120 minutes; aortic clamping time, considering prolonged when above 90 minutes; and mechanical circulatory assistance use (intra-aortic balloon pump).

Postoperative data

After surgery, patients were admitted to the Cardiovascular ICU still under anesthesia. They were intubated and mechanically ventilated (Evita 2 Dura® - Dräger Medical, Lübeck, Germany) in assisted/controlled volume mode,



* 1 due previous cardiac surgery, 37 submitted to combined surgery, 3 cause off-pump surgery, 1 due mini pump and 106 ventilated at admission with PEEP ≠ FiO₂ values different of 8 cmH₂O and/or 40% respectively.
 ** 2 without PaO₂ data, 3 due postoperative death e 2 for revision surgery.
 FiO₂: inspired oxygen fraction; PaO₂: partial pressure of oxygen in the arterial blood; PEEP: positive end-expiratory pressure.

Fig 1 - Patients admitted to the Cardiovascular Intensive Care Unit after coronary artery bypass graft

Table 1. Univariate analysis of preoperative variables. (n = 100).

Variables	Hypoxemia		P
	No (n = 45)	Yes (n = 55)	
Age (years)	62,6 ± 7,3	61,1 ± 11,5	0,453 ^a
BMI (kg/m ²)	24,9 ± 3,4	27,2 ± 3,8	0,002 ^a
Gender (%)			0,742 ^b
Female	26,7	21,8	
Male	73,3	78,2	
Hypertension			0,437 ^b
Absent	26,7	18,2	
Present	73,3	81,8	
DM			0,903 ^b
Absent	55,6	56,4	
Present	44,4	43,6	
Dyslipidemia			0,626 ^b
Absent	77,8	83,6	
Present	22,2	16,4	
AMI			0,673 ^b
Absent	82,2	87,3	
Present	17,8	12,7	
ARF			0,354 ^c
Absent	93,3	89,1	
Present	6,7	10,9	
Smoking			0,024 ^b
Absent	73,3	49,1	
Present	26,7	50,9	
COPD			0,388 ^c
Absent	97,8	94,5	
Present	2,2	5,5	
Angioplasty			0,575 ^c
Absent	97,8	96,4	
Present	2,2	3,6	

^aStudent's T Test ^bChi-square; ^cFisher Exact test ; BMI – Body Mass Index; DM – Diabetes Mellitus; ARF – Acute Renal Failure; AMI – Acute Myocardial Infarct; COPD – Chronic Obstructive Pulmonary Disease

according to Unit protocol [volume tidal (VT) 8 to 10 ml/kg; respiratory rate (RR) 12 to 16 bpm; inspitatory flow from 8 to 10 times minute volume (VT x RR); inspiratory time 1.0 second; FiO₂: 40%; PEEP: 8 cmH₂O].

Twenty minutes after admission, arterial blood was collected for a blood gas analysis (ABL 800 FLEX® - Radiometer, Bronshoj, Denmark), in which PaO₂ was analyzed and, consequently, gas exchange ratio (PaO₂/FiO₂). Other postoperative variables studied were mechanical ventilation duration and lengh of stay in the ICU.

According to 3rd Brazilian Consensus Conference on Mechanical Ventilation Consensus [5], normal PaO₂/FiO₂ ratio is above 300 mmHg; as gas exchange deterioration value equal or lower than 300 mmHg; and extreme severity of respiratory condition below 200 mmHg.

Current Berlin definition [10] classifies hypoxemia according to its level: Mild Hypoxemia (PaO₂/FiO₂ ≤ 300 and > 200 mmHg), Moderate Hypoxemia (PaO₂/FiO₂ ≤ 200 and >100 mmHg) and Severe Hypoxemia (PaO₂/FiO₂ ≤100 mmHg).

To determine the presence of gas exchange deterioration in researched subjects, it was considered hypoxemia a PaO₂/FiO₂ ratio ≤ 300, being classified according to Berlin definition [10].

Statistical analysis

Quantitative variables are described as mean and standard deviation or median (minimum, maximum), according to data distribution. Qualitative variables are presented by frequency and percentage. To normality test of the quantitative variables, D'Agostino-Pearson's test was used.

To test the relation between normally distributed quantitative variables and severe hypoxia occurrence, non-paired Student's t test was used. For quantitative variables of non-normal distribution, Mann-Whitney's test was used. Association between categorical and outcome variables was done using Chi-square test or Fisher Exact test. Quantitative variables whose statistic univariate tests presented P<0.25 were selected to enter the multivaried analysis by a multiple linear regression.

A 95%-confidence interval was set. To process and data analysis was used the software Bioestat version 5.3 (Instituto Mamirauá, Belém, Pará, Brazil).

RESULTS

The incidence of hypoxemia, i.e., in the presence of PaO₂/FiO₂ ≤300, in this study, was 55%.

About gender, most patients were male (76%). Mean age was 61.8±9.8 years (33-84 years) and 57% of them were elderly (≥ 60 years). There was no association between hypoxemia and these variables.

Mean BMI was 26.1±3.8 kg/m². Sixty percent of the sample were dystrophic. Among patients with hypoxemia in immediate postoperative period, 50.9% were overweight and

20% obese. This variable presented association to hypoxemia in this population (P=0.002).

Smoking was observed in 40% of patients. Among the patients who presented hypoxemia, 50.9% were smokers or former smokers. Mild level of COPD was prevalent in 4%. Hypertension prevailed in 78% of population and diabetes mellitus in 44%. Among comorbidities, smoking presents statistically association (P=0.024) for hypoxemia occurrence. Others variables were not statistically associated with development of hypoxemia in immediate postoperative period.

The univariate analysis of hypoxemia occurrence and preoperative variable of the study group are presented in Table 1.

Assessed intraoperative characteristics are shown in Table 2. Among studied patients, seven percent used only one graph, 40% two graphs and 53% three or more. Mean graphs number was 2.5±0.7, and LITA was used as graph in 91 patients. Remaining patients have received only venous graph. Average number of drains was 1.8±0.5. Twenty three percent used just one drain, seventy percent used two drains and 7% three drains. Mechanical circulatory assistance with intra-aortic balloon pump was used in only 3% of the subjects. None of these variables presented statistically association with hypoxemia in immediate postoperative period.

Mean surgery time was 3 hours and 45 minutes which was considered prolonged. Average pump time was 82.9±29 minutes. In 92% of the patients, pump time was shorter than 120 minutes. Mean aortic clamp time was 59.1±22.7, and 93% less than 90 minutes. These variables did not have any association with hypoxemia during CABG immediate postoperative.

For multiple regression, BMI and surgery time were considered. F value (regression) was significant (P=0.02), accepting the hypothesis that at least one of analyzed variables influenced hypoxemia occurrence. Among partial regression coefficients, only BMI was statistically significant (t=-2.66, P=0.009).

The median of time for interruption of ventilation device for the studied group lasted around 9 hours and 54 minutes and the ICU admission, two days. The presence of hypoxemia did not interfere with time of mechanical ventilation and ICU admission, as observed in Table 3.

Table 2. Univariate Analysis of the intra-operative variables (n= 100).

Variables	Hypoxemia		P
	NO (n= 45)	YES (n= 55)	
Number of grafts	2.5 ± 0.8	2.6 ± 0,7	0.537 ^a
LITA grafts (%)			0.354 ^b
Yes	6.7	10.9	
No	93.3	89.1	
Number of drains	1.8 ± 0.5	1.8 ± 0.5	0.934 ^a
Surgery time (min)	217.1 ± 48.1	231.6 ± 46.7	0.129 ^a
Pump time (min)	81.4 ± 29.8	84.1 ± 28.5	0.649 ^a
Aortic clamp time (min)	56.7 ± 23.2	61 ± 22.2	0.348 ^a

^a Student's T Test; ^b Fisher Exact Test; LITA – Left Internal Thoracic Artery

Table 3. Correlation between mechanical ventilation and permanence in the ICU, and hypoxemia time (N=100).

Variables	Hypoxemia		P
	NO (n = 45)	YES (n = 55)	
Mechanical ventilation duration (min)			
Median (Minimum; Maximum)	596 (88; 3872)	565 (120; 5425)	0,961 ^a
ICU length of stay (days)			
Median (Minimum; Maximum)	2 (2; 7)	2 (2; 7)	0,503 ^a

Mann Whitney; ICU - Intensive Care Unit

Mean PaO₂ quantified after ICU admission was 117.7±35.3 mmHg; in no hypoxemia patients was 150.6±22.2 mmHg; and in patients with hypoxemia was 72.1±3.9 mmHg. Average PaO₂/FiO₂ ratio was 294.2±88.2, ranging 165.3 to 510 mmHg.

When hypoxemia severity [10] was evaluated, 45% of 100 patients in the study had PaO₂/FiO₂ ratio classified as normal (> 300 mmHg). PaO₂/FiO₂ ratio greater than 200 mmHg and lower or equal to 300 mmHg (mild hypoxemia) was observed in 37% and 18% of them had moderate hypoxemia (100 mmHg<PaO₂/FiO₂<200 mmHg). None of the patients presented severe hypoxemia (PaO₂/FiO₂<100 mmHg).

DISCUSSION

Hypoxemia in CABG immediate postoperative is strictly related to risk factors associated to prior comorbidities and surgical procedure [11]. Among these comorbidities, advanced age, overweight, left ventricular dysfunction were associated to hypoxemia according to Szeles et al. [7], which analyzed risk factors to hypoxemia in CABG immediate postoperative in 481 patients with no relevant respiratory medical history.

In other studies [3,7,12], age was also mentioned as a risk factor related to hypoxemia after CABG. For Szeles et al. [7], age was an independent risk factor for hypoxemia and each year after age 34 increased their risk in 0.32% [7].

In our study we did not observe a significant association between advanced age and hypoxemia. However, considering physiological aspects, pulmonary function deteriorates with aging and the primary modification that happens in healthy person lungs due to aging is a gradual elastic retraction loss, reducing expiratory flow and increasing ventilation-perfusion inefficiency [13].

Divergent outcome from previous studies may be related to cut-off point of advanced age in this research, equal or above 60 years.

About weight, significant association was verified between high BMI and hypoxemia. This result is corroborated other studies in which obesity is described as a risk factor for hypoxemia [3,7].

Proposed mechanism to explain this result is obese patients' abdominal wall, in general, being thicker due to adipose panicle, contributing to increasing abdominal pressure over diaphragm and collapsed parenchyma volume in caudal and dependent lung areas [7].

Another associated factor with hypoxemia occurrence in this study was smoking. Ambrozin et al. [14] analyzed some pulmonary function aspects after CABG related to preoperative risks and verified that half of the cases presented decreased gas exchange ratio, influenced by comorbidities such as pre-existing lung disease, respiratory symptoms and smoking.

Smokers present higher respiratory complications risk during postoperative period, due not enough respiratory system integrity to maintain alveolar ventilation [15]. This data reinforces preoperative assessment importance to identify those patients [14].

According to Morsch et al. [16], damages in pulmonary function and respiratory muscle strength after CABG are associated to transoperative factors, such as sternotomy, CPB, and general anesthesia.

CPB has been considered important cause of hypoxemia in immediate postoperative CABG [3,7]. Szeles et al. [7] showed that CPB was an associated factor to severe hypoxemia in univariate analysis. Hypoxemia occurred in 37.9% of patients with pump time higher than 120 minutes; in 29.3% with pump time less than 120 minutes; and 20% in off-pump CABG.

In our research we did not observe an association between hypoxemia and CPB. This result may be related to lower pump time (less than 120 minutes) in 92% of patients, with mean time 82.9 minutes. According to Nozawa et al. [17], is highly probable that inflammatory aggression due longer pump time causes higher interstitial alveolar injury and edema.

However when Cox et al. [18] investigated effects of on and off-pump CABG on pulmonary function by an alveolar-arterial oxygen gradient in patients with adequate ventricular function and no pre-existing pulmonary disease, it was observed a similar level of pulmonary dysfunction in both groups. It suggests that pulmonary gas exchange deterioration associated with cardiac surgery is due to other factors, not only CPB. We highlight that in these patients, mean pump time was low (74.36 minutes), closer to our study time.

Szeles et al. [7] also demonstrated high incidence of hypoxemia (54.2%), even considering gas exchange lower than 200 mmHg. Among these, 27.3% presented PaO₂/FiO₂ ratio lower to 150, demonstrating higher hypoxemia severity in this group of patients when compared to our research, in which no one presented this ratio lower than 150 mmHg.

Some strategies to prevent hypoxemia have been used after cardio surgery, as alveolar recruitment with PEEP, showing gas exchange ratio improvement [19]. Almost half of patients needed some therapy to reverse hypoxemia such as alveolar recruitment, PEEP level elevation or oxygen therapy [20].

Laizo et al. [21] analyzed complications that increased ICU length of stay and found as cause those related to pulmonary function (COPD and smoking, pulmonary congestion, prolonged mechanical ventilation), besides infections, kidney failure, stroke and hemodynamic instability (as arterial hypotension, arrhythmia and AMI).

Weiss et al. [3] pointed hypoxemia as responsible for increase mechanical ventilation duration, ICU length of stay and hospital cost in CABG postoperative patients. This association was not observed in our research. Perhaps this result is explained by transitory hypoxemia in the first few hours after ICU admission presented by a lot of patients, with no implication in those times, besides PEEP level applied in admission.

Limitations of this study are related to its design, since retrospective studies may present selection and measurement biases. Regarding data collection from medical chart, with no direct patient's evolution follow-up may also be characterized as a limitation factor. In some included patients, variables related to lifestyle such as smoking habit and intensity, pre-existing diseases, and left ventricular function were only cited in a qualitative manner. While inevitable data fail, available variables were categorized in the best way possible.

CONCLUSION

In this sample, body mass index and smoking were associated to hypoxemia. These comorbidities are easy to be observed during preoperative assessment. For these high risk patients, implementation of mechanical ventilation strategies such as higher PEEP and alveolar recruitment, aiming to prevent and treat pulmonary collapse should contribute to minimize hypoxemia incidence and morbid-mortality related to cardiac surgery. Therefore, there are many aspects to study, yet, searching for effective interventions able to improve the clinical outcomes for this group of patients.

Author's roles & responsibilities	
NPS	Study design, data collection and manuscript writing
RMM	Study design and manuscript writing
DLB	Study design and manuscript writing
MAGC	Data collection
TEPB	Data collection
IML	Data collection
KCFM	Data collection
JLSL	Statistical analysis and text revision

REFERENCES

1. Piegas LS, Bittar OJNV, Haddad N. Cirurgia de revascularização miocárdica. Resultados do Sistema Único de Saúde. Arq Bras Cardiol. 2009;93(5):555-60.
2. Borghi-Silva A, Mendes RG, Costa FS, Di Lorenzo VA, Oliveira CR, Luzzi S. The influences of positive end expiratory pressure (PEEP) associated with physiotherapy intervention in phase I cardiac rehabilitation. Clinics (Sao Paulo). 2005;60(6):465-72.
3. Weiss YG, Merin G, Koganov E, Ribo A, Oppenheim-Eden A, Medalion B, et al. Postcardiopulmonary bypass hypoxemia: a prospective study on incidence, risk factors and clinical significance. J Cardiothorac Vasc Anesth. 2000;14(5):506-13.
4. Scanlan CL, Wilkins RL, Stoller JK. Fundamentos da terapia respiratória de Egan. 7ª ed. Barueri:Manole;2000.
5. Carvalho RRC, Toufen Júnior C, Franca AS. III Consenso de Ventilação Mecânica. Ventilação Mecânica: princípios, análise gráfica e modalidades ventilatórias. J Bras Pneumol. 2007;33(Suppl 2):S54-70.
6. Garutti I, Martinez G, Cruz P, Piñeiro P, Olmedilla L, la Gala F. The impact of lung recruitment on hemodynamics during one-lung ventilation. J Cardiothorac Vasc Anesth. 2009;23(4):506-8.
7. Szeles TF, Yoshinaga EM, Alencar W, Brudniewski M, Ferreira FS, Auler JO, et al. Hypoxemia after myocardial revascularization: analysis of risk factors. Rev Bras Anesthesiol. 2008;58(2):124-36.
8. Ji Q, Mei Y, Wang X, Feng J, Cai J, Sun Y, et al. Study on the risk factors of postoperative hypoxemia in patients undergoing coronary artery bypass grafting. Circ J. 2008;72(12):1975-80.
9. Organização Mundial da Saúde (OMS). Obesity: preventing and managing the global epidemic. Report of a WHO consultation on obesity. Geneva: WHO;1998.
10. The ARDS Definition Task Force. Acute respiratory distress syndrome: the Berlin definition. JAMA. 2012;307(23):2526-33.
11. Guizilini S, Gomes WJ, Faresin SM, Bolzan DW, Alves FA, Catani R, et al. Avaliação da função pulmonar em pacientes submetidos à cirurgia de revascularização do miocárdio com e sem circulação extracorpórea. Rev Bras Cir Cardiovasc. 2005;20(3):310-6.
12. Canver CC, Chanda J. Intraoperative and postoperative risk factors for respiratory failure after coronary bypass. Ann Thorac Surg. 2003;75(3):853-7.
13. Beliveau MM, Multach M. Perioperative care for the elderly patient. Med Clin North Am. 2003;87(1):273-89.
14. Ambrozini ARP, Caetano AJM. Aspectos da função pulmonar após revascularização do miocárdio relacionados com risco pré-operatório. Rev Bras Cir Cardiovasc. 2005;20(4):408-15.

-
15. Ferreira Júnior FH, Boas AGV, Gonnelli CA, Puig LB, Stolf N. Assistência ventilatória mecânica em pós-operatório de cirurgia cardíaca. *An Paul Med Cir.* 2003;130(1):22-30.
 16. Morsch KT, Leguisamo CP, Camargo MD, Coronel CC, Mattos W, Ortiz LDN, et al. Perfil ventilatório dos pacientes submetidos a cirurgia de revascularização do miocárdio. *Rev Bras Cir Cardiovasc.* 2009;24(2):180-7.
 17. Nozawa E, Kobayashi E, Matsumoto ME, Feltrim MI, Carmona MJ, Auler JJ. Assessment of factors that influence weaning from long-term mechanical ventilation after cardiac surgery. *Arq Bras Cardiol.* 2003;80(3):301-10.
 18. Cox CM, Ascione R, Cohen AM, Davies IM, Ryder IG, Angelini GD. Effect of cardiopulmonary bypass on pulmonary gas exchange: a prospective randomized study. *Ann Thorac Surg.* 2000;69(1):140-5.
 19. Auler JO Jr, Carmona MJ, Barbas CV, Saldiva PH, Malbouisson LM. The effects of positive end-expiratory pressure on respiratory system mechanics and hemodynamic in postoperative cardiac surgery patients. *Braz J Med Biol Res.* 2000;33(1):31-42.
 20. Vieira JE, Silva BAR, Garcia Júnior D. Padrões de ventilação em anestesia: estudo retrospectivo. *Rev Bras Anesthesiol.* 2002;52(6):756-63.
 21. Laizo A, Delgado FEF, Rocha GM. Complicações que aumentam o tempo de permanência na unidade de terapia intensiva na cirurgia cardíaca. *Rev Bras Cir Cardiovasc.* 2010;25(2):166-71.