

anger management. At the time of writing, the patient had not experienced any anger outbursts in the preceding 6 months and was in a stable relationship with a new girlfriend.

Table 1 summarizes first-line therapeutic approaches for IED.<sup>4</sup>

As illustrated in this case, IA is associated with substantial physical and psychosocial harm to the aggressive patient, the victims of the aggression, and society in general.<sup>4,5</sup> There is a need for better understanding of the role of IA in the high rates of violence observed in Brazil. As IA usually appears in adolescence or early adulthood and effective therapeutic approaches are available for its management,<sup>5</sup> proper treatment may significantly reduce personal and social distress.

Gustavo C. Medeiros,<sup>1</sup> Eric Leppink,<sup>2</sup> Liliana Seger,<sup>1</sup> Ana M. Costa,<sup>1</sup> Carolina Bernardo,<sup>1</sup> Hermano Tavares<sup>1</sup>  
<sup>1</sup>Department of Psychiatry, University of São Paulo, São Paulo, SP, Brazil. <sup>2</sup>Department of Psychiatry, University of Chicago, Chicago, IL, USA

Submitted Nov 08 2014, accepted Dec 03 2014.

## Disclosure

The authors report no conflicts of interest.

## References

- 1 Waiselfisz JJ. Mapa da violência 2014 [Internet]. In: Os jovens do Brasil. Brasília: FLACSO; 2014. 2014 [cited 2015 Jan 12]. [http://www.mapadaviolencia.org.br/pdf2014/Mapa2014\\_JovensBrasil.pdf](http://www.mapadaviolencia.org.br/pdf2014/Mapa2014_JovensBrasil.pdf).
- 2 Cleide Carvalho. Agressão física encabeça violência contra a mulher [Internet]. O Globo. 2014 Jun 15 [cited 2015 Jan 12]. <http://oglobo.globo.com/brasil/agressao-fisica-encabeça-violencia-contra-mulher-12866926#ixzz3ILtkQjas>.
- 3 Rogério Pagnan. PM separa 70 brigas de trânsito por dia [Internet]. Folha de São Paulo. 2011 Jul 10 [cited 2015 Jan 12]. <http://www1.folha.uol.com.br/multimedia/videocasts/941254-pm-separa-70-brigas-de-transito-por-dia-veja-relato-de-motorista-brigao.shtml>.
- 4 Coccaro EF, Lee R, McCloskey MS. Relationship between psychopathy, aggression, anger, impulsivity, and intermittent explosive disorder. *Aggress Behav*. 2014;40:526-36.
- 5 Coccaro EF. Intermittent explosive disorder as a disorder of impulsive aggression for DSM-5. *Am J Psychiatry*. 2012;169:577-88.

# Prevalence of psychiatric disorders among children and adolescents from four Brazilian regions

Rev Bras Psiquiatr 2015;37:178–179  
 doi:10.1590/1516-4446-2014-1606

Childhood psychiatric disorders usually persist through adolescence and adult life,<sup>1</sup> and have a deleterious impact and longstanding costs to individuals and society.<sup>2</sup> Ninety percent of the world's children and adolescents live in low- and middle-income countries, where methodologically sound data about the prevalence of child

psychiatric disorders are still lacking to guide improvement in services planning.<sup>1</sup>

In Latin America, studies examining the prevalence of psychiatric disorders in children using diagnostic instruments are very scarce. In Brazil, only three prevalence studies involving children's mental health and using diagnostic instruments have been conducted,<sup>3</sup> none of which involved more than one region of the country. Thus, the purpose of the current study was to ascertain the prevalence of psychiatric disorders in schoolchildren from grades 2-6 living in four municipalities from four Brazilian regions (Southeast, Center, Northeast, and North), using probabilistic samples.

Considering that education is compulsory for all Brazilians aged 4 to 17 years and that 83.5% of this population is enrolled in public schools, this multicenter, cross-sectional study enrolled 1,676 6-to-16-year-olds (response rate: 81.1%). Trained psychologists administered the Brazilian version of the Schedule for Affective Disorders and Schizophrenia for School-Age Children/Present-and-Lifetime-Version (K-SADS-PL) to mothers/main caregivers to identify current child psychiatric disorders. The Universidade de São Paulo Research Ethics Committee approved the study protocol. Data collection lasted 15 months, and was completed by December 2011. A detailed description of the methodology is available elsewhere.<sup>4</sup>

Prevalence rates were obtained for the overall sample and stratified by municipality/region, including any psychiatric disorder (excluding enuresis/encopresis), specific groups of disorders, and single disorders (Table 1). The observed overall prevalence of one or more psychiatric disorders (13.1%) was similar to most estimates reported for children around the world,<sup>2,5</sup> including rates from Brazil.<sup>3</sup> In comparison with one of the most important reviews in the field,<sup>5</sup> our estimates were similar in magnitude regarding any anxiety and any ADHD, somewhat smaller for any disruptive disorder, and smaller for any depressive and any oppositional conduct disorder.

Surprisingly, certain differences in prevalence rates by municipality/region were observed. The rate of any psychiatric disorder was lower in the Northeast municipality than in any of the other three sites. The rates of ADHD and disruptive disorders were higher in the Center of Brazil and lower in the Northeast, while oppositional/conduct disorder was more prevalent in the Center and less prevalent in the Northeast and North. These differences are difficult to interpret, as participating municipalities were similar in terms of population (having fewer than 50,000 inhabitants, as do 84.7% of Brazilian municipalities), were all located near a state capital, and had a Human Development Index near the countrywide average.

In conclusion, this was the first epidemiological study conducted in four Brazilian regions to investigate the prevalence of different types of psychiatric disorders among schoolchildren. Further studies are necessary to confirm or refute the differences observed by region. Because Brazil is a large and heterogeneous country in terms of socioeconomic status and availability of resources, local contexts must be examined to better capture differences in children's mental health needs.

**Table 1** Prevalence of psychiatric disorders in schoolchildren from four Brazilian regions (n=1,676)

Child psychiatric disorders	Overall sample (n=1,623)	Southeast Caeté (n=434)	Center Goianira (n=406)	Northeast Itaitinga (n=382)	North Rio Preto da Eva (n=401)	p
Any psychiatric disorder	13.1 (11.4-14.7)	14.5 (11.3-18.2)	18.5 (14.8-22.6)	6.8 (4.5-9.8)	12.0 (9.0-15.6)	<0.01
Any anxiety disorder	7.2 (6.0-8.8)	7.8 (5.5-10.8)	9.4 (6.7-12.6)	4.2 (2.4-6.7)	7.2 (4.9-10.2)	0.12
Separation anxiety disorder	1.3 (0.8-2.0)	2.3 (1.1-4.2)	1.5 (0.5-3.2)	0.8 (0.2-2.3)	0.7 (0.2-2.2)	0.15
Specific phobia	3.8 (2.9-4.9)	3.7 (2.1-5.9)	5.2 (3.2-7.8)	2.4 (1.1-4.4)	3.7 (2.1-6.1)	0.23
Social phobia	2.0 (1.4-2.8)	2.5 (1.3-4.5)	3.0 (1.5-5.1)	1.0 (0.3-2.7)	1.2 (0.4-2.9)	0.27
Posttraumatic stress disorder	0.3 (0.1-0.7)	0.5 (0.1-1.7)	0.2 (0.0-1.4)	0.0 (N/A)	0.5 (0.1-1.8)	0.57
Obsessive-compulsive disorder	0.6 (0.3-1.1)	0.2 (0.0-1.3)	0.2 (0.0-1.4)	0.8 (0.2-2.3)	1.0 (0.3-2.5)	0.38
Generalized anxiety disorder	0.4 (0.1-0.8)	0.5 (0.1-1.7)	0.7 (0.2-2.1)	0.0 (N/A)	0.2 (0.0-1.4)	0.37
Any depressive disorder	0.5 (0.2-1.0)	0.9 (0.3-2.3)	0.7 (0.2-2.1)	0.3 (0.0-1.4)	0.0 (N/A)	0.21
Major depression	0.2 (0.0-0.5)	0.5 (0.1-1.7)	0.2 (0.0-1.4)	0.0 (N/A)	0.0 (N/A)	0.34
Dysthymia	0.1 (0.0-0.4)	0.2 (0.0-1.3)	0.0 (N/A)	0.0 (N/A)	0.0 (N/A)	0.43
Any ADHD	4.5 (3.5-5.6)	4.6 (2.8-7.0)	8.1 (5.7-11.2)	1.0 (0.3-2.7)	4.0 (2.3-6.4)	<0.001
Any oppositional/conduct disorder	2.3 (1.6-3.1)	2.8 (1.4-4.8)	4.4 (2.6-6.9)	1.0 (0.3-2.7)	0.7 (0.2-2.2)	<0.001
Oppositional defiant disorder	1.7 (1.0-2.7)	2.5 (1.3-4.5)	3.4 (1.9-5.7)	0.3 (0.0-1.4)	0.5 (0.1-1.8)	<0.01
Conduct disorder	0.6 (0.3-1.1)	0.2 (0.0-1.3)	1.0 (0.3-2.5)	0.8 (0.2-2.3)	0.2 (0.0-1.4)	0.16
Any disruptive disorder*	5.8 (4.7-7.0)	6.2 (4.1-8.9)	9.9 (7.1-13.2)	2.1 (0.9-4.1)	4.7 (2.9-7.3)	<0.001
Other disorders†	1.5 (0.9-2.2)	2.3 (1.1-4.2)	1.5 (0.5-3.2)	1.0 (0.3-2.7)	1.2 (0.4-2.9)	0.04

Data presented as % (95%CI).

ADHD = attention deficit hyperactivity disorder; N/A = not applicable due to 0.0% prevalence.

\* Any disruptive disorder: any oppositional/conduct disorder and/or any ADHD.

† Other disorders: substance abuse, tic disorders, eating disorders, and/or psychotic disorders.

The next challenge is to translate evidence into action, scaling up services and human resources to deal appropriately with regional morbidity.

Cristiane S. Paula,<sup>1,2</sup> Evandro S. Coutinho,<sup>3</sup> Jair J. Mari,<sup>2</sup> Luis A. Rohde,<sup>4</sup> Euripedes C. Miguel,<sup>5</sup> Isabel A. Bordin<sup>2</sup>  
<sup>1</sup>Developmental Disorders Program, Universidade Presbiteriana Mackenzie, São Paulo, SP, Brazil. <sup>2</sup>Department of Psychiatry, Universidade Federal de São Paulo, São Paulo, SP, Brazil.  
<sup>3</sup>Department of Epidemiology, Fundação Oswaldo Cruz, Rio de Janeiro, RJ, Brazil. <sup>4</sup>Department of Psychiatry, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil. <sup>5</sup>Department of Psychiatry, Universidade de São Paulo, São Paulo, SP, Brazil.  
National Institute of Developmental Psychiatry for Children and Adolescents, São Paulo, SP, Brazil

Submitted Nov 03 2014, accepted Dec 10 2014.

### Acknowledgements

The study is part of the Instituto Nacional de Psiquiatria do Desenvolvimento para Infância e Adolescência (INPD), of the National Science and Technology Institute (Instituto Nacional de Ciência e Tecnologia, INCT). It was funded by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP; grant no. 2008/57896-8) and by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq; grant no. 573974/2008; Edital n° 15/2008).

The authors are grateful to Marcos Tomanik Mercadante (*in memoriam*) for his assistance with study design.

### Disclosure

The authors report no conflicts of interest.

### References

- Kieling C, Baker-Henningham H, Belfer M, Conti G, Ertem I, Omigbodun O, et al. Child and adolescent mental health worldwide: evidence for action. *Lancet*. 2011;378:1515-25.
- Merikangas KR, He JP, Brody D, Fisher PW, Bourdon K, Koretz DS. Prevalence and treatment of mental disorders among US children in the 2001-2004 NHANES. *Pediatrics*. 2010;125:75-81.
- Paula CS, Miranda CT, Bordin IAS. Saúde mental na infância e adolescência: revisão dos estudos epidemiológicos brasileiros. In: Lauridsen-Ribeiro E, Tanaka OY, editors. *Atenção em saúde mental para crianças e adolescentes no SUS*. São Paulo: Hucitec; 2010, p.75-92.
- Paula CS, Bordin IA, Mari JJ, Velasque L, Rohde LA, Coutinho ES. The mental health care gap among children and adolescents: data from an epidemiological survey from four Brazilian regions. *PloS One*. 2014;9:e88241.
- Costello EJ, Egger H, Angold A. 10-year research update review: the epidemiology of child and adolescent psychiatric disorders: I. Methods and public health burden. *J Am Acad Child Adolesc Psychiatry*. 2005;44:972-86.

## Psychotic syndrome secondary to meningioma treated with a low dose of olanzapine

Rev Bras Psiquiatr. 2015;37:179-180  
doi:10.1590/1516-4446-2014-1550

In November 2010, a 60-year-old woman presented for psychiatric evaluation. Her only words were: "It was not my fault." She remained silent for the rest of the interview, with a suspicious look. Her husband added that she had been very