



## Essential oils in the initial phase of broodstock diets of Nile tilapia

André Freccia<sup>1</sup>, Sílvia Maria de Negreiros Sousa<sup>1</sup>, Fábio Meurer<sup>2,4</sup>, Arno Juliano Butzge<sup>3,5</sup>,  
Juliana Kasper Mewes<sup>3,5</sup>, Robie Allan Bombardelli<sup>1,4</sup>

<sup>1</sup> Programa de Pós-Graduação em Zootecnia, Universidade Estadual do Oeste do Paraná, Marechal Cândido Rondon, PR, Brasil.

<sup>2</sup> Programa de Pós-Graduação em Aquicultura e Desenvolvimento Sustentável, Universidade Federal do Paraná, Palotina, PR, Brasil.

<sup>3</sup> Pontifícia Universidade Católica do Paraná, Toledo, PR, Brasil.

<sup>4</sup> Research Productivity Fellowship CNPq – Level 2.

<sup>5</sup> Scientific initiation fellowship.

**ABSTRACT** - This study evaluated the effect of feed supplementation (at doses of 0, 50, 100, 150, and 200 mg/kg) with SALUTO<sup>®</sup>, a microencapsulated blend of essential oils including carvacrol, cinnamaldehyde, 1,8-cineol, and pepper oleoresin, in Nile tilapia (*Oreochromis niloticus*) broodstock on reproductive and growth parameters during the initial phase of rearing. The growth parameters, somatic indexes and gonadal and hepatic parameters were analyzed. Quantitative parameters of growth, ovary fat and crude protein were not affected by supplementation of the additive. However, an increase in crude protein in the female liver was observed in correlation with increased essential oil levels. The female hepatosomatic index was also correlated with supplementation of essential oils. Feeding Nile tilapia broodstock diets containing an essential oil compound (SALUTO<sup>®</sup>) affects only the females, promoting an increase in liver protein inclusion and hepatosomatic index without impairing performance.

Key Words: additive, fish nutrition, growth promoter, hepatosomatic index

### Introduction

The accelerated growth of Brazilian aquaculture (MPA, 2010), associated with the intensification of fish farming, has resulted in animal health problems. The market pressure to reduce the use of antibiotics in aquaculture has been growing in recent years (Ardó et al., 2008). To reduce antibiotic resistance and avoid the presence of residues in food (Bruun et al., 2003), nutritional additives have been used in animal diets (Burt, 2004; Oncu et al., 2005).

The prohibition of growth promoters in Europe has stimulated the search for alternatives such as prebiotics (Silva and Nörnberg, 2003), probiotics (Santin et al., 2001; Meurer et al., 2006; Meurer et al., 2007; Meurer et al., 2008; Pozza et al., 2010), enzymes (Furuya et al., 2001; Surek et al., 2008), organic acids (Maiorka et al., 2004), and plant extracts (Çabuk et al., 2006; Santurio et al., 2007; Windisch et al., 2008; Applegate et al., 2010; Soltani et al., 2010). For example, specifically in fish, the use of garlic extract for silver catfish (*Rhamdia quelen*) (Fernandes et al., 2007) and propolis for the Nile Tilapia (Meurer et al., 2009a) has yielded satisfactory results.

The use of essential oils in the diet can promote improvement in the animal digestive tract (Chrubasik et al., 2005) by enhancing performance (Muhl and Liebert, 2007;

Santurio et al., 2007). These substances act by stimulating the secretion of digestive enzymes (Ramakrishnarao et al., 2003; Platel and Srinivasan, 2004), delaying gastric emptying (Manzanilla et al., 2004), accelerating the uptake of glucose by the intestine (Kreydiyyeh et al., 2003), or increasing the difficulty of pathogen adhesion to intestinal mucosa (Jamroz et al., 2006).

The initial phase of fish farming is important due to its influence on the other stages (Hayashi et al., 2002); appropriate growth is directly reflected in broodstock formation (Navarro et al., 2010). In the initial phase of broodstock formation, the use of additives that promote maximum growth (Traesel et al., 2011) can be an important management strategy.

There are many studies about administering essential oil blends to farm animals with contradictory results (Jesus, 2010; Barroca, 2011). However, a beneficial effect is observed when essential oils are used in situations of high yield production. Chilante et al. (2012) observed a reduction in the rate of cumulative mortality and increased egg production in broiler breeders using the same additive as that used in the present study.

In consideration of these observations, the use of essential oils could be an adequate and even strategic tool because it is a natural component without residual effects on

the meat, especially in Nile tilapia, the most important fish in the Brazilian aquaculture. However, no study regarding the use of essential oils in Nile tilapia was found.

The objective of this study was to evaluate the effects of increasing levels of a compound made from essential oils in reproductive and growth parameters of Nile tilapia (*Oreochromis niloticus*) broodstock diets during the initial phase of fish farming.

## Material and Methods

The experiment was conducted from January to June 2010 (180 days) in the Laboratory of Reproductive Technology of Cultivable Aquatic Animals (LATRAAC), located at Instituto de Pesquisa em Aquicultura Ambiental (InPAA), Universidade Estadual do Oeste do Paraná (UNIOESTE), Toledo Campus, Brazil. One thousand three hundred Nile tilapia (*Oreochromis niloticus*) from Thai lineage (13.66±0.16 g) stocked in 20 fine-mesh net cages called “hapas” (3 m × 2 m; 1 mm mesh) were used. Males and females were stocked at a density of 65 fish per hapa. The experiment was conducted in accordance with the ethical standards and approval of Colégio Brasileiro de Experimentação Animal (COBEA) and was approved by Comitê de Ética na Experimentação Animal e Aulas Práticas (CEEAAP/UNIOESTE) as protocol 06/10 minutes number 012012.

The hapas were installed in two earth ponds (20 × 10 m) and distributed in a completely randomized experimental design, with five treatments and four replicates. Treatments consisted of the provision of feed containing increasing levels of SALUTO<sup>®</sup>, a compound produced from various doses of essential oils (0, 50, 100, 150 and 200 mg/kg) as a microencapsulated blend of carvacrol, cinnamaldehyde, 1,8-cineol, and pepper oleoresin. One hapa containing 65 fish was considered as an experimental unit. Experimental diets containing 300 g/kg digestible protein (DP) and 3,000 kcal of digestible energy (DE)/kg of ration (Table 1) were formulated at the Laboratory of Aquatic Organisms Nutrition from the Technology in Aquaculture Graduate at Universidade Federal do Paraná, Palotina Sector, Brazil. To manufacture rations, crushed feed was pelleted in appropriate sizes according to the mouth of the fish (adapted from a protocol described by Bombardelli et al., 2009).

Fish count and measurements for each experimental parameter were performed every 15 days for the evaluation of growth and correction of feed rate. During this procedure, hapas were cleaned.

Fish were fed four times a day (8.00 h, 10.00 h, 14.00 h and 16.00 h) at a rate of 5.5% biomass. The feeding rate

was adjusted according to water temperature (adapted from Ostrensky and Boeger, 1998).

Water temperature was measured daily, in the morning and afternoon, with a mercury thermometer. Every two weeks, the levels of dissolved oxygen (YSI<sup>®</sup> 550A) and pH (TEC<sup>®</sup> TECNAL 5) of the water were measured at 6.00 h and 16.00 h (Reidel et al., 2010; Tessaro et al., 2012).

During the experimental period (January-June 2010), the mean water temperature was 21.63±3.67 °C (16.5 to 30 °C). The mean concentration of dissolved oxygen in the water during the experimental period was 4.28±1.55 mg/L in the morning and 5.54±1.10 mg/L in the afternoon. The mean values of water pH ranged from 7.02±0.36 in the morning to 7.03±0.22 in the afternoon. The recorded temperature, pH, and dissolved oxygen were suitable for the species, according to Hussain (2004) and Barbosa et al. (2007).

At the end of the experiment, the weight, standard length, average weight gain, daily weight gain, specific growth rate, apparent feed conversion ratio, survival rate (Bombardelli et al., 2009) and the condition factor (Vazzoler, 1996) were measured.

Afterwards, 80 fish (four females per experimental unit) were anesthetized and euthanized (CFMV, 2008). Liver, gonads, and viscera were weighed for the calculation of hepatosomatic, viscerosomatic (Bombardelli, 2007), and gonadosomatic indexes (Ng and Wang, 2011).

Table 1 - Composition of experimental diets with different levels of the compound made from essential oils for Nile tilapia broodstock diets


Ingredients (g/kg)	Treatments (mg/kg)				
	0	50	100	150	200
Soybean meal	553.100	553.100	553.100	553.100	553.100
Corn	317.300	317.300	317.300	317.300	317.300
Fishmeal	100.000	100.000	100.000	100.000	100.000
Supplement vit. + min. <sup>1</sup>	20.000	20.000	20.000	20.000	20.000
Salt (NaCl)	5.000	4.950	4.900	4.850	4.800
Dicalcium phosphate	4.300	4.300	4.300	4.300	4.300
Calcitic limestone	0.180	0.180	0.180	0.180	0.180
Propionic acid	0.100	0.100	0.100	0.100	0.100
Essential oil	0	50	100	150	200
Nutrients					
Linoleic acid (g/kg)	10.100	10.100	10.100	10.100	10.100
Starch (g/kg)	272.300	272.300	272.300	272.300	272.300
Ash (g/kg)	71.100	71.100	71.100	71.100	71.100
Fat (g/kg)	37.600	37.600	37.600	37.600	37.600
Digestible energy (kcal/kg)	3.000	3.000	3.000	3.000	3.000
Digestible protein (g/kg)	3.000	3.000	3.000	3.000	3.000
Crude fiber (g/kg)	28.300	28.300	28.300	28.300	28.300
Total lysine (g/kg)	18.400	18.400	18.400	18.400	18.400
Total phosphorus (g/kg)	10.000	10.000	10.000	10.000	10.000

<sup>1</sup> Supplement vitamin + mineral: composition of product in milligram per kilogram (Folic acid - 200 mg; pantothenic acid - 4,000 mg; biotin - 40 mg; copper - 2,000 mg; iron - 12,500 mg; iodine - 200 mg; manganese - 7,500 mg; niacin - 5,000 mg; selenium - 70 mg; vitamin A - 1,000,000 IU; vitamin B1 - 1,900 mg; vitamin B12 - 3,500 mg; vitamin B2 - 2,000 mg; vitamin B6 - 2,400 mg; vitamin C - 50,000 mg; vitamin D3 - 500,000 UI; vitamin E - 20,000 IU; vitamin K3 - 500 mg; zinc - 25,000 mg).

The livers from 40 females were fixed in Bouin aqueous solution for a period of 8 hours and transferred to a 70% alcohol solution. Then, the material was dehydrated by passage through a series of increasing concentrations of alcohol, diaphonized in xylene, and embedded in paraffin for obtaining semi-serial transverse sections with a thickness of 5  $\mu$ m. From each liver, two microscope slides were prepared containing five histological sections, which were stained with hematoxylin-eosin.

After histological processing, images were captured using a biological microscope at 40x magnification (Figure 1). In each histological section, five images were captured, totaling 25 images per liver. Measurements of average areas (Nikon microscope, model Eclipse 200, coupled to a Basler 602fc digital camera) of hepatocytes were performed according to Tessaro et al. (2012), using the software Image Pro-Plus 4.0.

Liver and gonads from the other 40 females were used for the analysis of crude protein from the gonads and livers and of the ether extract from the gonads. These analyses were performed in the Nutrition Laboratory at UNIOESTE – Marechal Cândido Rondon Campus, Brazil, according to Silva and Queiroz (2002).



A - hepatocyte; B - hepatocyte core; C - sinusoidal capillaries. Supplemented with essential oils at a dose of 200 mg/kg. Captured in biological microscope under 40x magnification.

Figure 1 - Liver tissue of female Nile tilapia.

The studied parameters were assessed by analysis of variance (ANOVA), and when differences were significant ( $P < 0.05$ ), the data were subjected regression analysis. All evaluations were performed using the software SAEG (Sistema para Análises Estatísticas e Genéticas, version 9.1).

## Results and Discussion

The parameters final weight, standard length, weight gain, daily weight gain, specific growth rate, apparent feed conversion, condition factor, and survival were not affected ( $P > 0.05$ ) by administration of the compound made from essential oils in the diets of the fish (Table 2).

Discussion of similar additives fed to fish is challenging because of the lack of experiments. However, comparisons with results from experiments with other species of farm animals, usually with monogastrics such as pigs and poultry (Chilante et al., 2012), are important. According to the results presented herein, additives fed to fish and other farming animals may not lead to increased performance parameters and may not provide different results in different species.

Using probiotics in the diet of Nile tilapia, Meurer et al. (2006, 2007, 2008, 2009b) found no effect on growth performance and survival. Meurer et al. (2009a), using brown propolis extract fed to Nile tilapia fingerlings, observed effects on growth and on feed conversion ratio. Barroca (2011) did not find effective results with SALUTO<sup>®</sup> fed to weaned piglets.

However, Jesus (2010) showed that SALUTO<sup>®</sup> improved weight gain, feed intake, feed conversion and production efficiency of broilers in the growth phase. Positive results were observed by Chilante et al. (2012), who fed a similar compound to broiler breeders using the same blend of essential oils.

Probiotics used in broilers showed a final weight and weight gain similar to observations in animals supplemented with antibiotic growth promoters (Çabuk et al., 2006, Traesel et al., 2011). Muhl and Liebert (2007) found that

Table 2 - Performance in the initial phase of broodstock rearing of Nile tilapia subjected to experimental diets with different levels of the compound composed of essential oils

Parameters	Treatments (mg/kg)					CV (%)
	0	50	100	150	200	
Average final weight (g)	80.73	80.05	82.05	80.97	81.48	8.26
Standard length (cm)	14.02	14.20	14.24	14.26	14.35	3.62
Average weight gain (g)	12.51	12.38	12.40	12.18	12.84	14.51
Average daily weight gain (g)	0.83	0.82	0.83	0.81	0.86	14.51
Specific growth rate (%)	0.47	0.48	0.48	0.47	0.49	7.09
Apparent feed conversion	1.54	1.55	1.59	1.58	1.51	8.54
Survival (%)	100.0	98.08	96.15	98.85	96.92	2.70

CV - coefficient of variation (%).

inclusion of a phytogetic additive in pig diets did not improve the performance or intestinal microflora of piglets. Hashemi and Davoodi (2010) and Hernandez et al. (2004) attributed the positive effects of plant extracts on growth performance and nutrient digestibility to the appetite and digestion-stimulating and antimicrobial properties of these compounds.

The addition of the essential oil compounds to the diet at levels above 50 to 200 mg/kg influenced ( $P < 0.05$ ) the female hepatosomatic index (Figure 2; Table 3) and increased crude protein inclusion in the female liver ( $P < 0.06$ ) (Figure 3; Table 5). On the other hand, the treatments did not affect ( $P > 0.05$ ) female gonadosomatic index, female viscerosomatic index, average area of female hepatocytes, female condition factor, male condition factor, male gonadosomatic index, male hepatosomatic index and male viscerosomatic index (Tables 3 and 4). The morphometry indexes of hepatocytes from female broodstocks were not affected by treatments ( $P > 0.05$ ) (Table 3).

In vertebrates, the liver is a key organ that controls many vital functions (Guyton and Hall, 2002). In fish, the liver plays an important role in females because during reproduction, this organ mobilizes food reserves for the production of vitellogenin (El-Sayed and Kawanna, 2007; Tsadik and Bart, 2007; Andrade et al., 2010). In this sense, the liver can be used as a body indicator of the nutritional or reproductive physiological status of fish (Caballero et al., 1999).

The increase in the female hepatosomatic index and liver crude protein might be a result of hyperplasia or hypertrophy (Caballero et al., 2004) due to greater liver overload derived from the diet. Ibrahim et al. (2011)

hypothesized that pancreatic protein enzymes trypsin and chymotrypsin produce hydrolyses and break down proteins into the basic amino acids.




Figure 2 - Effect of increasing levels of essential oils (mg/kg) in the diet of Nile tilapia broodstock on the female hepatosomatic index ( $P < 0.05$ ).




Figure 3 - Effect of increasing levels of essential oils (mg/kg) in the diet of Nile tilapia broodstock on the liver crude protein of females ( $P < 0.06$ ).

Table 3 - Somatic parameters and the area of hepatocytes in the initial phase of life of female Nile tilapia broodstock subjected to experimental diets with increasing levels of the essential oils compound

Parameters	Treatments (mg/kg)					CV (%)
	0	50	100	150	200	
Average area of female hepatocytes ( $\mu\text{m}^2$ )	127.24	135.58	135.27	124.76	126.03	20.26
Female gonadosomatic index (%)	3.73	3.94	3.65	4.08	4.14	23.71
Female viscerosomatic index (%)	11.95	11.59	12.10	12.23	12.40	9.27
Female condition factor	3.45	3.59	3.50	3.72	3.67	7.14

CV - coefficient of variation (%).

Table 4 - Somatic parameters in the initial phase of life of male Nile tilapia broodstock subjected to experimental diets with increasing levels of the essential oils compound

Parameters	Treatments (mg/kg)					CV (%)
	0	50	100	150	200	
Male gonadosomatic index (%)	0.76	0.76	1.15	0.96	0.82	32.88
Male hepatosomatic index (%)	2.16	2.35	2.71	2.34	2.44	16.12
Male viscerosomatic index (%)	9.17	9.74	10.71	10.02	10.12	13.95
Male condition factor	3.78	3.87	4.05	3.72	3.87	6.50

CV - coefficient of variation (%).

Table 5 - Chemical composition in the initial phase of life of female Nile tilapia subjected to experimental diets with different levels of the essential oil compound

Parameters	Treatments (mg/kg)					CV (%)
	0	50	100	150	200	
Crude protein in the female gonad (%)	42.85	31.65	41.91	32.51	26.20	30.69
Female gonadal fat (%)	7.99	6.33	8.71	4.13	8.17	59.27
Crude protein in the female liver <sup>1</sup> (%)	4.19	4.55	4.52	5.92	6.89	34.32

CV - coefficient of variation (%).

<sup>1</sup> Linear effect,  $y = 3.86164 + 0.0135304x$ ,  $r^2 = 0.86$ .

Some plants and bacteria stimulate trypsin secretion in the upper portion of the small intestine (Platel and Srinivasan, 2000), allowing a microflora modulating effect that might aid digestion and absorption of nutrients, thereby providing a greater array of amino acids for protein synthesis and thus increasing the body protein content.

As there were no changes in liver protein levels and hepatocyte area, the changes in hepatosomatic index are possibly related to the development of structures unrelated to the hepatocytes. Such structures might be related to the exocrine pancreatic tissue, or hepatopancreas, commonly present with diffuse distribution among the hepatic tissue of teleosts (Epple and Brinn, 1986), which is responsible for producing digestive enzymes that act in the intestine (Hinton and Pool, 1976; González et al., 1993).

In sex-reversed Nile tilapia fingerlings, the reduction of hepatosomatic index was observed when feed was supplemented with probiotic additives (Meurer et al., 2009a). Meurer et al. (2009b) suggested that this decrease in hepatosomatic index could be related to the nutritional aspect of fish and their growth rate, indicating that their metabolic reserves (glycogen/lipid) are used to enhance immunity against opportunistic organisms.

Because there was no health challenge in the experimental conditions, the fish used in this study might have directed their hepatic metabolism to produce vitellogenin and to promote the reproductive process. This process is most likely reflected in the liver metabolism and hepatosomatic index (Navarro et al., 2006). At the levels used in the present study, the essential oils compound did not elicit any toxicity problems. This is a positive factor because according to Traesel et al. (2011), when incorporated in excess, toxic effects of essential oils are observed. The most important mechanism of phytogetic feed additives arises from beneficially affecting the ecosystem of gut microflora through controlling potential pathogens (Hashemi and Davoodi, 2011).

The mode of absorption is a limiting factor for the use of feed additives for animals (Juven et al., 1994). The site of intestinal absorption of essential oils in monogastrics might influence its effect on the animal. Kohlert et al. (2000)

observed that the absorption of thymol in humans occurs in the upper intestine. In pigs, essential oils can be absorbed in the proximal portion of the intestine, and therefore, the antimicrobial activity in the distal portion of the intestine is limited (Muhl and Liebert, 2007). Thus, these factors may also influence the use of active principles present in the additive under study in this experiment because little information about the absorption modes of these substances through the digestive tract of tilapia is available.

Future experiments are needed to assess aspects of feeding essential oil compounds such as possible toxicity levels and the effects on microbiota and immunology at the various stages of development of farmed Nile tilapia.

## Conclusions

Feeding Nile tilapia broodstock diets containing an essential oil compound (SALUTO<sup>®</sup>) affects only the females, promoting increased liver protein inclusion and hepatosomatic index without diminishing performance.

## Acknowledgments

The authors thank Universidade Estadual do Oeste do Paraná, Universidade Federal do Paraná, GRASP Ind. e Com. Ltda., CNPq and CAPES.

## References

- Andrade, V. X. L.; Honji, R. M. and Romagoza, E. 2010. Processo de maturação das gônadas de pintado (*Pseudoplatystoma corruscans*) alimentado com dois níveis proteicos e suplementados com óleo de milho. Arquivo Brasileiro de Medicina Veterinária e Zootecnia 62:332-342.
- Applegate, T. J.; Klose, V.; Steiner, T.; Ganner, A. and Schatzmayr, G. 2010. Probiotics and phyto-genics for poultry: myth or reality? The Journal of Applied Poultry Research 19:194-210.
- Ardó, L.; Yin, G.; Xu, P.; Váradi, L.; Szigeti, G.; Jeney, Z. and Jeney, G. 2008. Chinese herbs (*Astragalus membranaceus* and *Lonicera japonica*) and boron enhance the non-specific immune response of Nile tilapia (*Oreochromis niloticus*) and resistance against *Aeromonas hydrophila*. Aquaculture 275:26-33.
- Barbosa, L. G.; Morais, G. and Inoue, L. A. K. A. 2007. Respostas metabólicas do matrinxã submetido a banhos anestésicos de eugenol. Acta Scientiarum Biological Sciences 29:255-260.

- Barroca, C. C. 2011. Aditivos em dietas para leitões de 21 a 49 dias de idade. Dissertação (M.Sc.). Universidade Federal de Viçosa, Viçosa, MG, Brasil.
- Bombardelli, R. A. 2007. Energia digestível para reprodutores de tilápia-do-nilo (*Oreochromis niloticus* L.). Tese (D.Sc.). Universidade Estadual de Maringá, Maringá.
- Bombardelli, R. A.; Hayashi, C.; Natali, M. R. M.; Sanches, E. A. and Piana, P. A. 2009. Desempenho reprodutivo e zootécnico e deposição de lipídios nos hepatócitos de fêmeas de tilápia-do-nilo alimentadas com rações de diversos níveis energéticos. Revista Brasileira de Zootecnia 38:1391-1399.
- Bruun, M. S.; Madsen, L. and Dalsgaard, I. 2003. Efficiency of oxytetracycline treatment in rainbow trout experimentally infected with *Flavobacterium psychrophilum* strains having different in vitro antibiotic susceptibilities. Aquaculture 215:11-20.
- Burt, S. 2004. Essential oils: their antibacterial properties and potential applications in foods – a review. International Journal of Food Microbiology 94:223-253.
- Caballero, M. J.; Izquierdo, M. S.; Kjorsvik, E.; Fernández, A. J. and Rosenlund, G. 2004. Histological alterations in the liver of sea bream, *Sparus aurata* L., caused by short- or long-term feeding with vegetable oils. Recovery of normal morphology after feeding fish oil as the sole lipid source. Journal of Fish Diseases 27:531-541.
- Caballero, M. J.; López-Calero, G.; Socorro, J.; Roo, F. J.; Izquierdo, M. S. and Fernández, A. J. 1999. Combined effect of lipid level and fish meal quality on liver histology of gilthead seabream (*Sparus aurata*). Aquaculture 179:277-290.
- Çabuk, M.; Bozkurt, M.; Alçiçek, A.; Akabağ, Y. and Kuçukyılmaz, K. 2006. Effect of a herbal essential oil mixture on growth and internal organ weight of broilers from young and old breeder flocks. South African Journal of Animal Science 36:135-141.
- CFMV - Conselho Federal de Medicina Veterinária. 2008. Resolução nº 876, de 15-02-2008. Publicada no DOU de 25-02-2008. Seção 1, p.100. Brasília, DF.
- Chilante, R. B.; Kussakawa, K. C. K. and Flemming, J. S. 2012. Efeitos da utilização de óleos essenciais na alimentas de aves matrizes pesadas. Revista Acadêmica Ciência Agrária Ambiental 10:387-394.
- Chrubasik, S.; Pittler, M. H. and Roufogalis, B. D. 2005. Zingiberis rhizoma: a comprehensive review on the ginger effect and efficacy profiles. Phytomedicine 12:684-701.
- El-Sayed, A. M. and Kawanna, M. 2007. Effects of photoperiod on growth and spawning efficiency of Nile tilapia (*Oreochromis niloticus* L.) broodstock in a recycling system. Aquaculture Research 38:1242-1247.
- Epple, A. and Brinn, J. E. 1986. Pancreatic islets. p.279-317. In: Vertebrate endocrinology: fundamental and biochemical implications. Pang, P. K. T. and Schereibman, M., eds. Academic Press, New York.
- Fernandes, M. C.; Garcia, J. R. E.; Muelbert, B. and Esquive, E. S. E. 2007. Utilização de extrato de alho na alimentação do jundiá *Rhamdia quelen* para controle de ictioftiriose. In: Anais da 2ª Jornada UNISUL de Iniciação Científica. UNISUL, Tubarão.
- Furuya, W. M.; Gonçalves, G. S.; Furuya, V. R. B. and Hayashi, C. 2001. Fitase na alimentação da tilápia do Nilo (*Oreochromis niloticus*). Desempenho e digestibilidade. Revista Brasileira de Zootecnia 30:924-929.
- González, G.; Crespo, S. and Brusle, J. 1993. Histo-cytological study of the liver of the cabrilla sea bass, *Serranus cabrilla* (Teleostei, Serranidae), an available model for marine fish experimental studies. Journal of Fish Biology 43:363-373.
- Guyton, A. C. and Hall, J. E. 2002. Tratado de fisiologia médica. 4.ed. Guanabara-Koogan, Rio de Janeiro.
- Hashemi, S. R. and Davoodi, H. 2010. Phytogenics as new class of feed additive in poultry industry. Journal of Animal and Veterinary Advances 9:2955-2304.
- Hashemi, S. R. and Davoodi, H. 2011. Herbal plants and their derivatives as growth and health promoters in animal nutrition. Veterinary Research Communications 35:169-180.
- Hayashi, C.; Boscolo, W. R.; Soares, C. M. and Meurer, F. 2002. Exigência de proteína digestível para larvas de tilápia do Nilo (*Oreochromis niloticus*), durante a reversão sexual. Revista Brasileira de Zootecnia 31:823-828.
- Hernandez, F.; Madrid, J.; Garcia, V.; Orengo, J. and Megias, M. D. 2004. Influence of two plant extracts on broiler performance, digestibility, and digestive organ size. Poultry Science 83:169-174.
- Hinton, D. E. and Pool, C. R. 1976. Ultrastructure of the liver in channel catfish *Ictalurus punctatus* (Rafinesque). Journal of Fish Biology 8:209-219.
- Hussain, M. G. 2004. Farming of tilapia: breeding plans, mass seed production and aquaculture techniques. Habiba Akter Hussain, Mymensingh.
- Ibrahim, S.H. A. M.; Omer, H. A. A.; Abedo, A. A.; Ali, F. A. F. and Abdel-Magid, S. S. 2011. Ginger root (*Zingiber officinale*) as feed additive in rabbit diets with two levels of protein. American-Eurasian Journal of Agricultural & Environmental Sciences 10:906-916.
- Jamroz, D.; Wiertelcki, T.; Houszka, M. and Kamel, C. 2006. Influence of diet type on the inclusion of plant origin active substances on morphological and histochemical characteristics of the stomach and jejunum walls in chicken. Journal of Animal Physiology and Animal Nutrition 90:255-268.
- Jesus, J. S. 2010. Utilização de probióticos, ácidos orgânicos e óleos essenciais na alimentação de frangos de corte. Dissertação (M.Sc.). Universidade Federal do Recôncavo da Bahia, Cruz das Almas.
- Juven, B. J.; Kanner, J.; Schved, F. and Weisslowicz, H. 1994. Factors that interact with the antibacterial action of thyme essential oil and its active constituents. Journal of Applied Bacteriology 76:626-631.
- Kohlert, C.; van Rensen, I.; März, R.; Schindler, G. and Veit, M. 2000. Bioavailability and pharmacokinetics of natural volatile terpenes in animals and humans. Planta Medica 66:495-505.
- Kreydiyyeh, S. I.; Usta, J.; Knio, K.; Markossian, S. and Dagher S. 2003. Aniseed oil increases glucose absorption and reduces urine output in the rat. Life Sciences 74:663-673.
- Maiorka, A.; Santin, A. M. E.; Borges, S. A.; Opalinski, M. and Silva A. V. F. 2004. Emprego de uma mistura de ácidos fumárico, láctico, cítrico e ascórbico em dietas iniciais de frangos de corte. Archives of Veterinary Science 9:31-37.
- Manzanilla, E. G.; Perez, J. F.; Martin, M.; Kamel, C.; Baucells, F. and Gasa, J. 2004. Effect of plant extracts and formic acid on the intestinal equilibrium of early-weaned pigs. Journal of Animal Science 82:3210-3218.
- Meurer, F.; Hayashi, C.; Costa, M.M.; Mauerwerk, V.L. and Freccia, A. 2006. Utilização de *Saccharomyces cerevisiae* como probiótico para tilápias-do-nilo durante o período de reversão sexual submetidas a um desafio sanitário. Revista Brasileira de Zootecnia 35:1881-1886.
- Meurer, F.; Hayashi, C.; Costa, M.M.; Freccia, A. and Mauerwerk, M.T. 2007. *Saccharomyces cerevisiae* como probiótico para alevinos de tilápia-do-Nilo submetidos a um desafio sanitário. Revista Brasileira de Zootecnia 36:1219-1224.
- Meurer, F.; Hayashi, C.; Costa, M.M.; Mascioli, A. S.; Saragiotto, L. M. and Freccia, A. 2008. Levedura como probiótico na reversão sexual da tilápia-do-Nilo. Revista Brasileira de Saúde e Produção Animal 9:804-812.
- Meurer, F.; Costa, M. M.; Barros, D. A. D.; Oliveira, S. T. L. and Paixão, P. S. 2009a. Brown propolis extract in feed as a growth promoter of Nile tilapia (*Oreochromis niloticus*, Linnaeus 1758) fingerlings. Aquaculture Research 40:603-608.
- Meurer, F.; Silva, M. S.; Costa, M. M.; Colpini, L. M. S. and Mascioli, A. H. 2009b. Probiótico com levedura na alimentação da tilápia do Nilo, durante o período de reversão sexual, cultivada em água de tanque de cultivo. Revista Brasileira de Saúde e Produção Animal 10:406-416.
- MPA - Ministério da Pesca e Aquicultura. 2010. Produção de pescado aumenta 25%. Available at: <[http://www.mpa.gov.br/#imprensa/2010/AGOSTO/nt\\_AGO\\_19-08-Producao-de-pescado-aumenta](http://www.mpa.gov.br/#imprensa/2010/AGOSTO/nt_AGO_19-08-Producao-de-pescado-aumenta)> Accessed on: Sep. 25, 2011.

- Muhl, A. and Liebert, F. 2007. Growth and parameters of microflora in intestinal and faecal samples of piglets due to application of a phytogetic feed additive. *Journal of Animal Physiology and Animal Nutrition* 91:411-418.
- Navarro, R. D.; Matta, S. L. P.; Lanna, E. A. T.; Donzele, J. L.; Rodrigues, S. S.; Silva, R. F.; Calado, L. L. and Ribeiro Filho, O. P. 2006. Níveis de energia digestível na dieta de piauçu (*Leporinus macrocephalus*) no desenvolvimento testicular em estágio pós-larval. *Zootecnia Tropical* 24:153-163.
- Navarro, R. D.; Maldonado, I. R. S. C.; Matta, S. L. P.; Ribeiro Filho, O. P.; Pereira, F. K. S.; Rodrigues, S. S. and Calado, L. L. 2010. Associação do nível de energia digestível no comprimento total, peso das gônadas e índice gonadosomático de fêmeas de Piauçu ("*Leporinus macrocephalus*", SPIX 1829) em estágio pós-larval. *Revista Brasileira de Saúde e Produção Animal* 11:242-251.
- Ng, W-K and Wang, Y. 2011. Inclusion of crude palm oil in the broodstock diets of female Nile tilapia, *Oreochromis niloticus*, resulted in enhanced reproductive performance compared to broodfish fed diets with added fish oil or linseed oil. *Aquaculture* 314:122-131.
- Oncu, S.; Erdem, H. A. and Pahsa, A. 2005. Therapeutic options for pneumococcal pneumonia in Turkey. *Clinical Therapeutics* 27:674-683.
- Ostrensky, A. and Boeger, W. 1998. Piscicultura: fundamentos e técnicas de manejo. Agropecuária, Guaíba.
- Platel, K. and Srinivasan, K. 2000. Influence of dietary spices and their active principles on pancreatic digestive enzymes in albino rats. *Nahrung* 44:42-46.
- Platel, K. and Srinivasan, K. 2004. Digestive stimulant action of spices: a myth or reality? *Indian Journal of Medical Research* 119:167-179.
- Pozza, P. C.; Pozza, M. S. S.; Nunes, R. V.; Pasquetti, T. J.; Venturi, I. and Busanello, M. 2010. Desempenho, microbiota intestinal e peso de órgãos de leitões na fase inicial recebendo rações com simbiótico e probiótico. *Ciência e Agrotecnologia* 34:1327-1334.
- Ramakrishnarao, R.; Platel, K. and Srinivasan, K. 2003. In vitro influence of spices and spice-active principles on digestive enzymes of rat pancreas and small intestine. *Nahrung* 47:408-412.
- Reidel, A.; Boscolo, W. R.; Feiden, A. and Romagosa, E. 2010. The effect of diets with different levels of protein and energy on the process of final maturation of the gametes of *Rhamdia quelen* stocked in cages. *Aquaculture* 298:354-359.
- Santin, E.; Maiorka, A. and Macari, M. 2001. Performance and intestinal mucosa development of broiler chickens fed diets containing *Sacharomyces cerevisiae* cell wall. *The Journal of Applied Poultry Research* 10:236-244.
- Santurio, J. M.; Santurio, D. F.; Pozzatti, P.; Moraes, C.; Franchin, P. R. and Alves, S. H. 2007. Atividade antimicrobiana dos óleos essenciais de orégano, tomilho e canela frente a sorovares de *Salmonella enterica* de origem avícola. *Ciência Rural* 37:803-808.
- Silva, L. P. and Nörnberg, J. L. 2003. Prebióticos na nutrição de não ruminantes. *Ciência Rural* 33:983-990.
- Silva, D. J. and Queiroz, A. C. 2002. Análise de alimentos: métodos químicos e biológicos. 3.ed. Editora UFV, Viçosa, MG, Brasil.
- Soltani, M.; Sheikhzadeh, N.; Ebrahimzadeh-Mousavi, H. A. and Zargar, A. 2010. Effects of *Zataria multiflora* Essential Oil on innate immune responses of common carp (*Cyprinus carpio*). *Journal of Fisheries and Aquatic Science* 5:191-199.
- Surek, D.; Maiorka, A.; Dahlke, F.; Opalinski, M.; Franco, S. G. and Krabbe, E. L. 2008. Uso de fitase em dietas de diferentes granulometrias para frangos de corte na fase inicial. *Ciência Rural* 38:1725-1729.
- Tessaro, L.; Toledo, C. P. R.; Neumann, J.; Krause, R. A.; Meurer, F.; Natali, M. R. M. and Bombardelli, R. A. 2012. Growth and reproductive characteristics of *Rhamdia quelen* males fed on different digestible energy levels in the reproductive phase. *Aquaculture* 326-329:74-80.
- Traesel, C. K.; Wolkmer, P.; Schmidt, C.; Silva, C. B.; Paim, F. C.; Rosa, A. P.; Alves, S. H.; Santurio, J. M. and Lopes, S. T. A. 2011. Serum biochemical profile and performance of broiler chickens fed diets containing essential oils and pepper. *Comparative Clinical Pathology* 20:453-460.
- Tsadik, G. G. and Bart, A. N. 2007. Effects of feeding, stocking density and water-flow rate on fecundity, spawning frequency and egg quality of Nile tilapia, *Oreochromis niloticus* (L.). *Aquaculture* 272:380-388.
- Vazzoler, A. E. A. M. 1996. Biologia da reprodução de peixes teleósteos: teoria e prática. EDUEM, Maringá.
- Windisch, W.; Schedle, K.; Plitzner, C. and Kroismayr, A. 2008. Use of phytogetic products as feed additives for swine and poultry. *Journal of Animal Science* 86(E. Suppl.):E140-E148.