



A new species of *Trichodesmium* (Cyanobacteria) from freshwaters, Brazil

Célia L. Sant'Anna^{1,3}, Watson A. Gama¹, Thais G. Silva² & Armando A.H. Vieira²

Abstract

Trichodesmium is a typical planktic genus in which trichomes are disposed either in fascicles or radially in rounded colonies. Based on morphological studies, there are eleven species of *Trichodesmium*, out of which nine are from marine environments and only two from freshwaters. *Trichodesmium* is mainly known for its capacity to form blooms and produce toxins in marine tropical and subtropical environments. There is no information about the capacity of the freshwater *Trichodesmium* species to produce toxins. It was only with molecular studies that the taxonomy of marine *Trichodesmium* started to be solved. However, up to now, no material has been available for molecular analyses of freshwater species. During the studies of microalgae from São Paulo state, a population resembling *Trichodesmium* was found in a recreational pond. The analyzed organisms formed fascicles of homocyted and not attenuated trichomes and cells with gas vesicles, a set of features that makes them different from the other freshwater *Trichodesmium* species. Thus, we have described the species *Trichodesmium brasiliense* sp. nov. based on material from Brazilian inland water. Also, we have suggested revision of some Brazilian literature citations of *T. lacustre* and their inclusion in the synonym of this new species.

Key words: new species, Oscillatoriales, shallow lakes, *Trichodesmium lacustre*, tropical region.

Resumo

Trichodesmium é um gênero tipicamente planctônico cujos tricomas estão dispostos em fascículos ou radialmente em colônias. Com base em estudos morfológicos, existem 11 espécies de *Trichodesmium* descritas, das quais nove são para ambientes marinhos e apenas duas para águas continentais. Este gênero é conhecido principalmente por sua capacidade de formar florações e produzir toxinas em ambientes marinhos tropicais e subtropicais. Contudo, não há informações sobre produção de toxinas por suas espécies de água-doce. Foi apenas com os estudos moleculares que a taxonomia das espécies marinhas começou a ser resolvida. Contudo, até o presente, não há material disponível para a realização de estudos moleculares com as espécies de ambientes continentais. Durante o estudo das microalgas do Estado de São Paulo, uma população similar a *Trichodesmium* foi encontrada em uma lagoa utilizada para recreação. Os organismos analisados formavam fascículos compostos por tricomas homocitados e não atenuados, com aerótopos presentes; características que os tornam diferentes de outras espécies de *Trichodesmium*. Assim, descrevemos a espécie *Trichodesmium brasiliense* sp. nov. a partir da análise de espécimes de águas continentais brasileiras. Adicionalmente, sugerimos a revisão de algumas citações da literatura brasileira para *T. lacustre* e inclusão dessas na sinonímia da nova espécie descrita.

Palavras-chave: nova espécie, Oscillatoriales, lagos rasos, *Trichodesmium lacustre*, regiões tropical/subtropical.

Introduction

Trichodesmium was originally proposed by Ehrenberg (1830) and later was included in the genus *Oscillatoria* (Kützing 1843). It was only based on the work of Gomont (1892) that the genus *Trichodesmium* was definitively considered as a valid generic entity. According to Komárek & Anagnostidis (2005), *Trichodesmium* is typically

planktic, in that the trichomes are disposed either in fascicles or radially in rounded colonies and are cylindrical or slightly attenuated, straight or curved, rarely coiled, without sheath and always present gas vesicles. Based on morphological studies, Komárek & Anagnostidis (2005) recognized 11 species of *Trichodesmium*, out of which nine are from marine environments and only two are from freshwaters.

¹ Instituto de Botânica, Lab. Ficologia, Av. Miguel Stéfano 3687, 04301-012, São Paulo, SP, Brazil

² Universidade Federal de São Carlos, Depto. Botânica, Rod. Washington Luís s/n, km 235, 13565-905, São Carlos, SP, Brazil

³ Author for correspondence: celialsant@gmail.com

Trichodesmium is mainly known for its capacity for forming blooms and producing toxins in marine tropical and subtropical environments (Capone *et al.* 1997; Orcutt *et al.* 2002; Detoni *et al.* 2016). According to Suvapepun (1992), the blooms are initially formed in deep layers of the water column and later they accumulate on the water surface forming an extended reddish or yellowish mass. The ability to accumulate on the water surface is due to the big and very resistant gas vesicles of the *Trichodesmium* genus (Capone *et al.* 1997).

Trichodesmium species play a very important role in nitrogen fixation and, in some marine environments, they constitute most part of the total biomass. Biomass contributes intensively to nitrogen fixation and, consequently, contributes significantly to the global nitrogen cycle (Carpenter 1983; Capone *et al.* 1997; Capone *et al.* 2005).

In relation to toxin production, the marine *Trichodesmium* species are known as microcystin and saxitoxin producers (Jackson *et al.* 2001; Shaw *et al.* 2001; Long & Carmichael 2003). In the Brazilian literature, there are also records of intoxication induced by *Trichodesmium* blooms in marine environments (Detoni *et al.* 2016), but there is no information of the capacity of the freshwater *Trichodesmium* species to produce toxins.

The taxonomy of marine *Trichodesmium* started to be really solved only with molecular studies (Orcutt *et al.* 2002). According to these studies, there are only two groups of marine species: one formed exclusively by *Trichodesmium erythraeum* Ehrenb. ex Gomont and the other formed by closely related species (*Trichodesmium thiebautii* Gomont ex Gomont, *Trichodesmium tenue* Wille, *Trichodesmium hildebrandtii* Gomont and *Katagnymene spiralis* Lemmerm.). There is no molecular information about freshwater *Trichodesmium* but in terms of phylogeny, a great distance between the marine and freshwater species will probably be found on the generic level (Komárek & Anagnostidis 2005). Unfortunately, up to now there is no material for the necessary molecular analyses of freshwater species.

The two freshwater species, *Trichodesmium iwanoffianum* Nygaard and *Trichodesmium lacustre* Kleb., have been poorly studied and rarely cited in the literature. The first is restricted to tropical regions and the second occurs in temperate, tropical and subtropical areas (Komárek & Anagnostidis 2005). There are few references about freshwater

Trichodesmium in Brazil and all of them refer to *T. lacustre* (Komárek & Komárková-Legnerová 2007; Martins *et al.* 2012). However, the species has been problematic since its description, when Klebahn pointed out its similarity to heterocyst-free *Aphanizomenon* representatives (Klebahn 1895). Despite its confirmation as a separate taxon and identity, Prescott (1962) also commented that the material he found in freshwater environments in North America may not exactly correspond to *T. lacustre*, although it was the closest species to his material. This puts in evidence that the diversity of continental *Trichodesmium* species is greater than it has been described so far.

A population similar to *Trichodesmium*, with homocyted trichomes disposed in fascicles and cells with gas vesicles, was found during studies on microalgae from São Paulo state. The organisms present a set of morphological features that makes them different from the other *Trichodesmium* species. Thus, based on our results, we are proposing a new species of *Trichodesmium* from Brazilian inland water.

Material and Methods

The studied sample was collected with a plankton net (20 µm) in a shallow pond used for recreation, coordinates 21° 03' 302 S & 050° 02' 809 W. The sample was preserved in formaldehyde 4%. The type material is kept in the Herbarium of Institute of Botany, São Paulo, Brazil, under the accession number SP469314. Sample aliquots of analyzed material by Martins *et al.* (2012) and Komárek & Komárková-Legnerová (2007) were recovered from Herbaria to compare the population described by these authors with the herein proposed species.

Based on nature material, the morphological study was performed with a Zeiss Axioplan 2 optical microscope and the identification and cell measurements were carried out in 15 individuals. Digital photographs taken with a Zeiss Axiocam MRc digital camera documented the organisms. The observed morphological features were the following: fascicle length and width and cell length, width and length/width rate, presence of gas vesicle and trichome features (attenuation, constriction and form of apical cell). The new species was described according to the International Code of nomenclature for algae, fungi, and plants (McNeill *et al.* 2012) and the classification system proposed by Komárek *et al.* (2014) was adopted.

Results and Discussion

Trichodesmium brasiliense Sant'Anna et al.

Fig. 1a-j

Synonymous: *Trichodesmium lacustre sensu* Martins et al. (2012) and *Trichodesmium cf. lacustre sensu* Komárek & Komárková-Legnerová (2007).

Thallus microscopic, fasciculate, 70–96 μm wide, 820–868.5 μm long, formed by several (6–10) trichomes densely aggregated. Trichomes straight, not attenuated, strongly constricted, without mucilage. Cells elongated or more rarely barrel shaped, with gas vesicles, 4.6–7.1 μm wide, 4.9–7.4 μm long; apical cell hemispheric or elongated.

Holotype: BRAZIL. SÃO PAULO: Zacarias, Prainha do Zacarias, 19.III.2014, T.G. Silva (SP469314).

Habitat: shallow pond, presence of macrophytes (*Eichhornia*, *Salvinia*, *Egeria* and *Eleocharis*); pH 7.15, water temperature 32 °C, dissolved oxygen 3.81 mg L⁻¹; conductivity 0.023 mS cm⁻¹.

Table 1 shows that there are evident morphological differences between the compared species: the studied population presents deeply constricted trichomes and rounded-elongated and isodiametric cells (width/length rate around 1) different from *Trichodesmium iwanoffianum*, which has short barrel-shaped cells (width/length rate 2–3.5). As for *T. lacustre*, the greatest difference is the attenuated trichomes of the




Figure 1 – *Trichodesmium brasiliense* – a-b. general aspect of fascicles; c-d. details of apex of fascicles; e-g. details of trichomes with rounded cells, gas vesicles and granules; h-i. details of smaller trichomes with elongated cells and uniform content; j. hormogonium.

Table 1 – Comparison between the Brazilian material and the other *Trichodesmium* species from freshwaters.

<i>Trichodesmium</i>	Fascicle dimensions (μm)	Trichomes	Form of cells	Cell dimensions (μm)	Apical cell	Distribution
<i>T. iwanoffianum</i> Nygaard	50–100 \times 650–900	Straight or slightly curved, cylindrical, constricted	Short barrel	7–8 \times 2–5	Hemispherical	Tropical: Sumatra, Mozambique, Caribbean, China, Brazil
<i>T. lacustre</i> Klebahn	-----	Straight, attenuated, constricted	Short barrel	5–8 \times 3–6	Cylindrical with rounded apex, rounded, or hemispheric	Temperate, tropical: Europe, USA, Burma, India, China.
<i>T. brasiliense</i> sp. nov.	70–96 \times 868.5	Straight, cylindrical, constricted	Elongated to barrel-shaped	4.6–7.1 \times 4.9–7.4	Hemispheric to elongated	Tropical and subtropical: Brazil

species in contrast to the cylindrical trichomes of *T. brasiliense*. Length/width rate together with trichome width are very important morphologic markers to the group of homocyted cyanobacteria, mainly for the Oscillatoriaceae and Microcoleaceae families (Komárek *et al.* 2014). Besides, the Brazilian material presents two types of trichomes inside the same fascicle: Figure 1e-g shows strongly constricted trichomes with rounded-elongated cells and Figure 1h-i displays smaller and less constricted trichomes with barrel-shaped cells and some of them without gas vesicles and granules. These two types of trichomes in the same fascicle have never been described to other freshwater species. Considering that both types of trichomes are densely aggregated in the fascicle, it is possible to suppose that the smaller trichomes with some cells without gas vesicles and granules (“lighter”) would be involved in nitrogen fixation, exactly as described by Bergman *et al.* (2013). According to these authors, this process is known to marine species of *Trichodesmium* to perform spatial fixation of nitrogen, which is much more efficient than temporal fixation.

Based on the statement of Komárek & Anagnostidis (2005), the material mentioned by Smith (1918, 1920) as *Trichodesmium lacustre* does not correspond to the original description of this species because it does not present attenuated trichomes. Thus, Komárek & Anagnostidis (2005) analyzed and identified the material just as *Trichodesmium* sp. Comparing the Brazilian population with the literature data, and with the additional material recovered from Brazilian Herbaria, we conclude that the specimens presented by Martins *et al.* (2012) as *T. lacustre* to South Brazil and those from Komárek & Komárková-Legnerová (2007) referred as *T. cf lacustre* to São

Paulo state, are identical to our studied population and should be resigned into *Trichodesmium brasiliense*. As we could see, these materials also do not present attenuated trichomes, what is a quite common feature to *T. lacustre*. Also in the literature, the populations showed by Smith (1918, 1920) as *Trichodesmium lacustre* and by Tiffany (1937) and Prescott (1962) as *Oscillatoria lacustris* (Kleb.) Geitler, all of them from USA lakes, morphologically match the Brazilian material and certainly belong to the same species. However, as we have only morphological data obtained from literature and the ecology and geographic distance are different from Brazilian populations, we just highlight that these USA specimens could also belong to *T. brasiliense*. It is interesting that Prescott (1962) already pointed out his doubt about the identification of his material: “The specimens assigned here are enigmatic. The bundles of trichomes strongly suggest *Trichodesmium* because of their arrangement and lack of heterocysts. They are not like *Trichodesmium lacustre* Klebahn, however, in respect to the morphology of the apical cell which in that species is long and attenuated, but in our specimens it is short and rotund. There is scarcely enough evidence at present to justify giving a new name to the Wisconsin plant, but it is possible that subsequent study will establish it as a new freshwater species of *Trichodesmium*”. According to Komárek & Komárková-Legnerová (2007) “the identity of all described populations over the world up to now (including the Brazilian ones) should be revised”.

The description of new species of cyanobacteria without genetic data is not very usual in current days. However, the accessibility to molecular information is not the same to all cyanobacterial groups, due to restrictions regarding

pure culturing and single cell isolation from nature populations. So, in the cases of particular evidences, as morphoecological markers that distinguish populations which cannot be accessed by molecular techniques, the description of new taxa works as the recognition of the unknown biodiversity, which is proved to be vast to cyanobacteria (Nabout *et al.* 2013). *T. brasiliense* fits in this case by its singular morphological features, and differently from marine environments, in freshwater the populations of *Trichodesmium* showed to be very small. Summarizing, we are describing a new species of *Trichodesmium*, named *T. brasiliense*, which occurs in lakes from South America, encompassing tropical and subtropical regions.

Acknowledgments

The authors want to thank FAPESP (process 2011/50054-4) for the financial support and WAG also thanks FAPESP for fellowship support (FAPESP 2012/16430-1). We are also deeply thankful to Dr. Jiří Komárek, from the Academy of Science of the Czech Republic, for his excellent comments on freshwater *Trichodesmium*.

References

- Bergman B, Sandh G, Lin S, Larsson J & Carpenter EJ (2013) *Trichodesmium*—a widespread marine cyanobacterium with unusual nitrogen fixation properties. *FEMS microbiology reviews* 37: 286-302.
- Capone DG, Burns JA, Montoya JP, Subramaniam A, Mahaffey C, Gunderson T, Michaels AF & Carpenter EJ (2005) Nitrogen fixation by *Trichodesmium* spp.: An important source of new nitrogen to the tropical and subtropical North Atlantic Ocean. *Global Biogeochemical Cycles* 19.
- Capone DG, Zehr JP, Paerl HW, Bergman B & Carpenter EJ (1997) *Trichodesmium*, a globally significant marine cyanobacterium. *Science* 276: 1221-29.
- Carpenter EJ (1983) Nitrogen fixation by marine *Oscillatoria*. *Nitrogen in the marine environment*: 65-104.
- Detoni AMS, Costa LDF, Pacheco LA & Yunes JS (2016) Toxic *Trichodesmium* bloom occurrence in the southwestern South Atlantic Ocean. *Toxicon* 110: 51-55.
- Ehrenberg C (1830) Neue Beobachtungen über blutartige Erscheinungen in Aegypten, Arabien und Sibirien, nebst einer Uebersicht und Kritik der früher bekannten. *Annalen der Physik* 94: 477-514.
- Gomont MA (1892) Monographie des Oscillariées (Nostocacées homocystées). *Annales des Sciences Naturelles, Botanique, Série 7* 15: 263-368, pls 6-14.
- Jackson W, Carmichael W & Carpenter E (2001) The paralytic shellfish toxins, N-sulfocarbamoyl and gonyautoxin, are the major neurotoxic factors associated with the bloom-forming *Trichodesmium* spp. Abstracts, Fifth International Conference on Toxic Cyanobacteria, Noosa Lakes.
- Klebahn H (1895) Gasvacuolen, ein Bestandteil der Zellen der wasserblütbildenden Phycochromaceen. *Flora* 80: 241-82, pl. IV.
- Komárek J & Anagnostidis K (2005) Cyanoprokaryota 2. Teil Oscillatoriales. In: Büdel B, Krienitz L, Gärtner G & Schagerl M (eds.) Süßwasserflora von Mitteleuropa. Elsevier Spektrum Akademischer Verlag, München. P. 759.
- Komárek J, Kaštovský J, Mareš J & Johansen, J.R. 2014. Taxonomic classification of cyanoprokaryotes (cyanobacterial genera) 2014, using a polyphasic approach. *Preslia* 86: 295-335.
- Komárek, J. & Komárková-Legnerová, J. 2007. Several rare freshwater planktic Cyanobacteria (Cyanoprokaryotes) from reservoirs in South America. *Hoehnea* 34: 49-58.
- Kützing FT (1843) *Phycologia generalis: oder Anatomie, Physiologie und Systemkunde der Tange*. Brockhaus,
- Long B & Carmichael W (2003) Marine cyanobacterial toxins. In: Hallegraeff GM, Anderson DM, Cembella AD & Enevoldsen HO (eds.) *Manual on harmful marine microalgae*. Unesco, Paris. Pp. 279-96.
- Martins MD, Branco LHZ & Werner VR (2012) Cyanobacteria from coastal lagoons in southern Brazil: non-heterocytous filamentous organisms. *Brazilian journal of botany* 35: 325-38.
- McNeill J, Barrie F, Buck W, Demoulin V, Greuter W, Hawksworth D, Herendeen P, Knapp S, Marhold K & Prado J (2012) *International code of nomenclature for algae, fungi and plants. Regnum vegetabile*. Vol. 154. Koeltz Scientific Books, Koenigstein. 240p
- Nabout JC, Silva Rocha B, Carneiro FM & Sant'Anna CL (2013) How many species of Cyanobacteria are there? Using a discovery curve to predict the species number. *Biodiversity and conservation* 22: 2907-18.
- Orcutt K, Rasmussen U, Webb EA, Waterbury JB, Gundersen K & Bergman B (2002) Characterization of *Trichodesmium* spp. by genetic techniques. *Applied and environmental microbiology* 68: 2236-45.
- Prescott GW (1962) *Algae of the western Great Lakes area*. 2nd ed. Wm.C. Brown, Dubuque. 977 p.
- Shaw G, Poon K, Codd G, O'Neil J, Lam P, Chong M, Davis B, Eaglesham G & Moore M (2001) The

- presence of microcystin-like compounds in blooms of the marine cyanobacterium, *Trichodesmium* spp. Fifth International Conference on Toxic Cyanobacteria, Noosa. CSIRO, Canberra. P. 1.
- Smith GM (1918) A second list of algae found in Wisconsin lakes. Transactions of the Wisconsin Academy of Science, Arts and Letters 19: 614-54.
- Smith GM (1920) Phytoplankton of the inland lakes of Wisconsin. Part I: Myxophyceae, Phaeophyceae, Heterokontae, and Chlorophyceae exclusive of the Desmidiaceae. Bulletin of the Wisconsin Geological and Natural History Survey 57: 1-243.
- Suvapepun S (1992) *Trichodesmium* blooms in the Gulf of Thailand. Marine pelagic cyanobacteria: *Trichodesmium* and other diazotrophs. Springer, Dordrecht. Pp. 343-48.
- Tiffany LH (1937) The filamentous algae of the west end of Lake Erie. American Midland Naturalist 18: 911-51.

Editora de área: Dra. Renata Reis

Artigo recebido em 05/09/2017. Aceito para publicação em 22/11/2017.



This is an open-access article distributed under the terms of the Creative Commons Attribution License.