

# Sex-differential effects of olanzapine vs. aripiprazole on glucose and lipid metabolism in first-episode schizophrenia

XUE-MEI ZHOU<sup>1</sup>  
<http://orcid.org/0000-0002-4774-4432>

MAO-RONG HU<sup>2</sup>  
<http://orcid.org/0000-0002-8541-2893>

MEI-YU GONG<sup>1</sup>  
<http://orcid.org/0000-0002-9092-3006>

XUE-LIANG ZOU<sup>1</sup>  
<http://orcid.org/0000-0003-0688-1550>

ZHI-MIN YU<sup>1</sup>  
<http://orcid.org/0000-0002-8732-4811>

<sup>1</sup> Jiangxi Mental Health Centre, Nanchang, Jiangxi, China.

<sup>2</sup> The First Affiliated Hospital of Nanchang University, Nanchang, Jiangxi, China.

Zhou XM et al. / Arch Clin Psychiatry. 2019;46(2):33-9

Received: 06/29/2018 – Accepted: 04/02/2019

DOI: 10.1590/0101-60830000000189

On Table 4, where is written:

**Table 4.** The between-drug and within-drug evolution of the individual metabolic parameter over time in male patients with first-episode schizophrenia assigned to 12 weeks treatment with aripiprazole or olanzapine

	NBMI	NTG
Baseline, female	0.41 (-0.05–0.88)	-0.66 (-1.24–0.09)
(The 4th weekend - Baseline), female	0.17 (0.01–0.32)	0.70 (0.22–1.18)
(The 8th weekend - Baseline), female	0.26 (0.11–0.42)	0.95 (0.47–1.43)
(The 12th weekend - Baseline), female	0.47 (0.32–0.63)	1.21 (0.73–1.69)
Baseline × (male - female)	-0.40 (-1.06–0.26)	0.21 (-0.60–1.03)
The 4th weekend × (male - female)	0.08 (-0.13–0.30)	0 (-0.68–0.68)
The 8th weekend × (male - female)	0.21 (-0.01–0.42)	-0.07 (-0.74–0.61)
The 12th weekend × (male - female)	0.16 (-0.06–0.38)	0.14 (-0.54–0.82)

Data are effect size (95% CI). The measurement units of all laboratory values for the fasting serum levels of lipids and glucose use mmol/L. All indicators are shown by their respective normal score. BMI: body mass index; TG: triglyceride.

It should be read:

**Table 4.** The between-drug and within-drug evolution of the individual metabolic parameter over time in male patients with first-episode schizophrenia assigned to 12 weeks treatment with aripiprazole or olanzapine

	NBMI	NTG
Baseline, aripiprazole	0.41 (-0.05–0.88)	-0.66 (-1.24–0.09)
(The 4th weekend - Baseline), aripiprazole	0.17 (0.01–0.32)	0.70 (0.22–1.18)
(The 8th weekend - Baseline), aripiprazole	0.26 (0.11–0.42)	0.95 (0.47–1.43)
(The 12th weekend - Baseline), aripiprazole	0.47 (0.32–0.63)	1.21 (0.73–1.69)
Baseline × (olanzapine - aripiprazole)	-0.40 (-1.06–0.26)	0.21 (-0.60–1.03)
The 4th weekend × (olanzapine - aripiprazole)	0.08 (-0.13–0.30)	0 (-0.68–0.68)
The 8th weekend × (olanzapine - aripiprazole)	0.21 (-0.01–0.42)	-0.07 (-0.74–0.61)
The 12th weekend × (olanzapine - aripiprazole)	0.16 (-0.06–0.38)	0.14 (-0.54–0.82)

Data are effect size (95% CI). The measurement units of all laboratory values for the fasting serum levels of lipids and glucose use mmol/L. All indicators are shown by their respective normal score. BMI: body mass index; TG: triglyceride.

DOI: 10.1590/0101-60830000000200

